

The *Rhizobium*-Plant Symbiosis

PIETERNEL VAN RHIJN† AND JOS VANDERLEYDEN*

F. A. Janssens Laboratory of Genetics, KU Leuven, B-3001 Heverlee, Belgium

INTRODUCTION	124
Diversity of Legumes	124
Taxonomy and Host Specificity of <i>Rhizobium</i> Species	124
Infection and Nodulation Mechanisms	125
Rhizobial Genes Controlling Infection, Nodulation, and Host Range: an Overview	127
STRUCTURAL <i>nod</i> GENES AS HOST RANGE DETERMINANTS	127
Common <i>nod</i> Genes	127
Host-Specific <i>nod</i> Genes	127
REGULATORY <i>nod</i> GENES AS HOST RANGE DETERMINANTS	129
<i>nodD</i> Gene	129
NodD as a member of the LysR family of transcriptional activators	129
NodD as a membrane protein	129
Plant signals activating NodD	130
NodD as a determinant of host range	130
NodD regulation	131
Other <i>nod</i> Regulatory Genes	131
Nod FACTORS	133
Biological Activities of Nod Metabolites	133
Biochemical Function of the <i>nod</i> Gene Products	133
Common <i>nod</i> genes and the molecular backbone	133
Synthesis of specific side chains	134
Nod Metabolites: Variation on a Theme	135
ROLE OF NodO IN SYMBIOSIS	135
CONCLUSION	136
ACKNOWLEDGMENTS	136
REFERENCES	136

INTRODUCTION

Under conditions of nitrogen limitation, *Rhizobium*, *Bradyrhizobium*, and *Azorhizobium*, collectively referred to as rhizobia, elicit on their leguminous hosts the formation of specialized organs, nodules. In these root or stem structures, the bacteria are able to convert atmospheric nitrogen into ammonia, which is used by the plant as a nitrogen source.

Diversity of Legumes

The ability to establish a nitrogen-fixing symbiosis with rhizobia is restricted to legumes, with one exception, the genus *Parasponia* of the Ulmaceae (262). The Leguminosae family comprises three subfamilies, Caesalpinioideae, Mimosoideae, and Papilionoideae, each of which contains genera able to form root nodules (2, 190). In these three subfamilies, the percentages of nodulated species are quite different. There are just a few nonnodulating genera in the advanced subfamilies Papilionoideae and the Mimosoideae, but the less specialized subfamily Caesalpinioideae includes many nonnodulating genera (45). Since the Caesalpinioideae is the most primitive subfamily of the Fabales, it is assumed that the symbiosis was

developed at a relatively late stage during legume evolution (288). This notion is also supported by the fact that several *Parasponia* species in the family Ulmaceae form legume-like nitrogen-fixing nodules with a variety of rhizobial strains which can also nodulate some legumes. Since the members of the Ulmaceae are no more closely related to the members of the Leguminosae than to almost any other family of dicotyledons, no real phylogenetic sense can be made of this phenomenon (288). Leguminous plants are very diverse in morphology, habitat, and ecology, ranging from Arctic annuals to tropical trees (45). Because a large number of legumes are nodulated by rhizobia, the symbiosis with rhizobia is apparently not an adaptation to a specialized ecological niche but, rather, depends on some genetic peculiarity of legumes, one that is so complex that it has rarely evolved elsewhere in the plant kingdom (288).

Taxonomy and Host Specificity of *Rhizobium* Species

The three rhizobial genera, *Rhizobium*, *Bradyrhizobium*, and *Azorhizobium*, have for many years been grouped with the agrobacteria and phyllobacteria into one family, the *Rhizobiaceae* (127). The use of modern methods of bacterial systematics, such as numerical taxonomy, nucleic acid hybridization, and 16S rRNA analysis, has demonstrated the existence of marked genetic diversity within this family (287). It is now widely accepted that *Rhizobium* and *Bradyrhizobium* are only distantly related (167). Each of these genera has close relatives that are not plant symbionts and are placed in different families. This is most evident for *Bradyrhizobium* (283, 285, 287).

The current taxonomic classification of the rhizobia is given

* Corresponding author. Mailing address: F. A. Janssens Laboratory of Genetics, KU Leuven, W. de Croylaan 42, B-3001 Heverlee, Belgium. Phone: 32 16 32 2396. Fax: 32 16 32 2990. Electronic mail address: Anita.vermassen@agr.kuleuven.ac.be.

† Present address: Department of Biology, University of California Los Angeles, Los Angeles, CA 90024-1606.

TABLE 1. Rhizobium-plant associations

Rhizobium	Host plant(s)
<i>R. meliloti</i>	<i>Medicago</i> , <i>Melilotus</i> , and <i>Trigonella</i> spp.
<i>R. leguminosarum</i> bv. <i>viciae</i>	<i>Pisum</i> , <i>Vicia</i> , <i>Lathyrus</i> , and <i>Lens</i> spp.
bv. <i>trifolii</i>	<i>Trifolium</i> spp.
bv. <i>phaseoli</i>	<i>Phaseolus vulgaris</i>
<i>R. loti</i>	<i>Lotus</i> spp.
<i>R. huakuii</i>	<i>Astragalus sinicus</i>
<i>R. ciceri</i>	<i>Cicer arietinum</i>
<i>Rhizobium</i> sp. strain NGR234.....	Tropical legumes, <i>Parasponia</i> spp. (nonlegume)
<i>R. tropici</i>	<i>Phaseolus vulgaris</i> , <i>Leucaena</i> spp., <i>Macroptilium</i> spp.
<i>R. etli</i>	<i>Phaseolus vulgaris</i>
<i>R. galegae</i>	<i>Galega officinalis</i> , <i>G.</i> <i>orientalis</i>
<i>R. fredii</i>	<i>Glycine max</i> , <i>G. soja</i> , and other legumes
<i>B. japonicum</i>	<i>Glycine max</i> , <i>G. soja</i> , and other legumes
<i>B. elkanii</i>	<i>Glycine max</i> , <i>G. soja</i> , and other legumes
<i>Bradyrhizobium</i> sp. strain <i>Parasponia</i>	<i>Parasponia</i> spp.
<i>A. caulinodans</i>	<i>Sesbania</i> spp. (stem nodulating)

in Table 1. The species name of the microsymbionts reflects in most cases the corresponding host plant nodulated and suggests that symbiosis is a species-specific process. The situation is much more complex than can be reflected in Table 1, in which some of the host plants are matched up with the microsymbiont. It is quite clear that the degree of host specificity varies tremendously among the rhizobia (288). Some strains have a very narrow host range, for example *Rhizobium leguminosarum* bv. *trifolii*, while others, like *Rhizobium* sp. strain NGR234, have a very broad host range. It is becoming increasingly clear that the symbiotic relationships between legume species and rhizobia is quite complex, as illustrated by the following examples. *Rhizobium* sp. strain NGR234 nodulates at least 35 different legume genera as well as the nonlegume *Parasponia* (153). Recently, it was demonstrated that *Azorhizobium caulinodans* can nodulate *Phaseolus vulgaris* (280). According to the data in Table 1, rhizobia isolated from alfalfa should belong to the species *R. meliloti*. However, Eardly et al. (67) have shown that some rhizobia isolated from alfalfa, including strain OR191, can nodulate *Phaseolus vulgaris*. Classification analysis of the 16S ribosomal sequence of OR191 have shown that strain OR191 is more closely related to *R. etli* and other bean symbionts than to *R. meliloti* (167). Unfortunately, this isolate was not further characterized with respect to the genes involved in nodulation. With the identification, cloning, and mutagenesis of the nodulation genes (see below), it has been shown that inactivation of a single gene drastically alters the host range of a particular strain (71, 118). *R. leguminosarum* bv. *viciae* and *R. leguminosarum* bv. *trifolii* are very similar with respect to their symbiotic genes. However, their host ranges are very different. Exchange of one gene in the microsymbiont can switch the host range of the two biovars (246). Plant tests to determine the nodulation phenotype are nevertheless important, as they allow verification of predicted host range on the basis of characterization of the molecular structure of the signals that govern nodulation.

Infection and Nodulation Mechanisms

Root infection by rhizobia is a multistep process that is initiated by preinfection events in the rhizosphere. Rhizobia respond by positive chemotaxis to plant root exudates and move toward localized sites on the legume roots (11, 28, 56, 79, 98). Both *Bradyrhizobium* and *Rhizobium* spp. are attracted by amino acids, dicarboxylic acids present in the exudates, and very low concentrations of excreted components, such as flavonoids, that may not have high nutritional value (1, 4, 27, 129, 187). Apparently, chemotaxis is not required for nodulation, because flagellum-deficient mutants are still able to nodulate normally; instead, it has an influence on competition and the establishment in the rhizosphere (158, 163, 171).

Subsequently, the rhizobia attach to the plant root surface. For many rhizobia, primary target sites for infection are young growing root hairs, but there are no exclusive loci for rhizobial attachment (238). Initially it was speculated that attachment of *Rhizobium* and *Bradyrhizobium* spp. to legume roots was involved in host specificity. Specific adherence of compatible rhizobia was proposed to be mediated by specific binding of particular polysaccharide structures present on the bacterial cell surface to host plant lectins (19, 102). The lectin recognition hypothesis has been validated in one case in which clover plants transgenic for a pea lectin gene can be nodulated by *R. leguminosarum* bv. *viciae*, which normally does not nodulate clover (53a). Many studies failed to demonstrate any degree of host specificity at the attachment step (174, 193, 277). Smit et al. (233–237) undertook a detailed study on the attachment capacity of *R. leguminosarum* bv. *viciae* to pea root hairs under various physiological conditions. It was concluded that the conditions under which the rhizobia were grown strongly influenced the attachment to pea roots. Under specific growth conditions, lectin seems to be clearly involved in attachment of rhizobia (133). Smit et al. (234) demonstrated that rhizobial attachment is a two-step process, at least in the *Rhizobium*-pea system. In the first step, *R. leguminosarum* binds loosely as single cells to the root hair surface. In the second step, referred to as cap formation, additional bacteria accumulate at the adhesion site. However, it is still unclear which determinants are involved in each of these two steps. Under laboratory conditions, the firm attachment step is not even essential for nodulation (278). For *R. leguminosarum* bv. *viciae*, a Ca^{2+} -dependent adhesin, called rhicadhesin, mediates the initial direct attachment to pea root hair surfaces (234, 239). For cap formation, the firm attachment step, fibrilous appendages of (brady)rhizobia appear to be involved. These appendages can be cellulose fibrils (*R. leguminosarum*) or proteinous fimbriae (*Bradyrhizobium japonicum*) (234, 277, 279). However, it cannot be excluded that other bacterial macromolecules might be involved as well (110, 111).

At the surface of the root, and very probably also from a distance, rhizobia cause root hair branching, deforming, and curling (16, 269, 286). These phenomena are also observed with supernatants of induced rhizobia cultures. In all *Rhizobium*-plant interactions studied thus far, the active substances in the supernatant have been identified as lipooligosaccharides, also called Nod factors. These Nod factors are synthesized by means of some of the nodulation genes (150, 245, 263). In legumes, the region that is most susceptible to *Rhizobium* infection is just behind the apical meristem at the site of emerging root hairs (15). Elongated root hairs are sometimes infected, but this is rather unusual (29, 194). Mostly, the young root hairs can be curled sufficiently to entrap bacterial cells in a pocket of host cell wall. After entrapment, a local lesion of the root hair cell wall is formed by hydrolysis of the cell wall

(10, 179, 180, 265). The mechanism of hydrolysis of the cell wall is not known. Either the bacteria may induce hydrolytic enzymes that are responsible for localized cell wall dissolution, or the bacteria may exploit plant mechanisms such as those used when epidermal cells grow out into root hairs (132). Rhizobia enter the roots at the sites where root hair cell walls are hydrolyzed. The penetration occurs by invagination of the plasma membrane. The host plant reacts by depositing new cell wall material around the lesion in the form of an inwardly growing tube (29, 266). The tube is filled with proliferating bacteria surrounded by a matrix and becomes an infection thread. The infection thread grows toward the inner tangential wall of the root hair cell tip by a process of tip growth (10).

Concomitantly with formation of the infection thread, particular cortical cells divide to form a nodule primordium, and the infection thread grows toward these primordia (156, 181, 273, 284). In soybean (30, 69) and alfalfa (63, 284), cell division in the cortex can also occur prior to the invasion of the root hairs. In *Glycine max*, the first divisions occur in the hypodermis and spread to the outer layers of the cortex (30). Most division centers are not clearly associated with curled root hairs or (later) hairs with infection threads; most of these cell division centers are arrested and never form nodules (30). The root cortical cells through which infection threads will pass on their way to the nodule primordia change markedly before they are penetrated by an infection thread. Detailed cytological analyses have shown that microtubuli rearrange, that the nucleus migrates to the cell center, and that additional cell wall material is formed. On the basis of these cytological changes, it has been suggested that the cortical cells become prepared for infection thread penetration (267).

The location of the nodule primordia in the root cortex depends on the type of nodule formed by a particular plant (181). In general, in temperate legumes such as pea, vetch, and alfalfa, the primordium is formed from cells in the inner cortex (63, 156). These legumes form indeterminate cylindrical nodules and have a persistent apical meristem (180). This persistent activity of the meristem ensures nodule elongation, since new cells are constantly added to the distal end of the nodule (272). While the meristem is active, rhizobia are released from the infection threads into the plant cell cytoplasm (22, 108, 131, 181). The differentiation of micro- and macrosymbiont leads to the establishment of a central zone of the nodule, in which nitrogen is reduced (177, 272). Thus, in indeterminate nodules, nodule growth and functioning occur simultaneously, and all intermediates in differentiation can be observed in a single longitudinal section of a nodule. On the other hand, in most tropical legumes, such as soybean and French bean, nodules have a determinate growth pattern (179, 266). A nodule meristem is induced in the root outer cortex, and the bacteria are released into actively dividing meristematic cells, each daughter cell receiving rhizobia (179, 181). Meristematic activity is restricted to a short period. Following a round of successive divisions, the invaded meristematic cells differentiate simultaneously to form the nitrogen-fixing central tissue (181). As a result of this developmental pathway, nodule growth and function are dissociated. Determinate nodules do not elongate but enlarge, and only a single stage of plant and bacterial differentiation can be observed at any particular moment.

Besides the formation of infection threads through root hairs, which is most widely studied, the rhizobia may enter through cracks in the epidermis. In legumes such as *Arachis hypogaea* (peanut) and *Stylosanthes* spp., microsymbionts infect their hosts by "crack entry" (32, 33). In the presence of rhizobia, cell divisions are induced in the cortex of an emerging lateral root. Growth of the young root causes separation of

cortical and epidermal cells and enables entry and intercellular spread of rhizobia. In both genera, no infection threads are formed, and rhizobia colonize the root apoplast presumably by cell wall digestion or, in *Stylosanthes* spp., by progressive collapse of outer root cells. Continuous host cell divisions result in development of a uniformly infected central tissue resembling the determinate nodule type (32, 33, 250).

Also, stem and root nodules of *Sesbania rostrata* are induced following crack entry by *A. caulinodans* at the base of dormant root primordia, which are present in rows along the length of the stem, or at the base of secondary roots in the case of root nodules (62, 65, 178, 264). Direct intercellular infection is followed by very active multiplication of the bacteria, forming wide intercellular spaces filled with azorhizobia. These spaces then extend inward as narrow, branched, intercellular infection threads which spread into the meristematic zone induced in the cortex. The subsequent release of bacteria into the cytoplasm of newly induced meristematic cells leads eventually to the development of a determinate nodule (65, 178, 263).

Bradyrhizobium sp. infects *Parasponia andersonii* (Ulma-ceae), the only nonlegume plant genus to form nitrogen-fixing root nodules with rhizobia (261). Upon inoculation, the first sign of root nodule initiation is the formation of swollen multicellular root hairs. Simultaneously, the colonizing bacteria stimulate cell division in the outer cortex. In due course, these cell divisions cause development of callus-like bumps which rupture the epidermis, especially at the base of multicellular root hairs, and infection follows through these wounds (crack entry). After invasion, a large proliferation of intercellular rhizobia occurs, which is associated with damage to the host cells. Occasionally, intracellular rhizobia in single file surrounded by plant cell wall material are observed. Rhizobia are not endocytotically released from these infection threads, but the infection thread will change in nature as rhizobia differentiate into the nitrogen-fixing bacteroid form (147, 260, 261). This type of interaction is generally considered a primitive rhizobial infection, since the orderly invasion of the intercellular space (which leads to the formation of fixing threads), and the more damaging pathogenic type of interaction can occur in relatively close proximity (147). This type of invasion and infection thread formation has been described for root nodules of many genera in the family Caesalpinioideae and some in the family Papilionoideae, like *Andira* (46, 47).

A third mode of infection is observed in *Mimosa scabrella*, a tropical tree, where the rhizobial infection sites are at junctions of epidermal cells (44). The bacteria penetrate the radial walls and proliferate intercellularly. Occasionally, subepidermal root hairs were formed upon inoculation, but they were never infected. Penetration of outer root cells starts from irregular cell wall ingrowth and progresses in the cortical region through the primary wall layer, rather than by separating cells at the middle lamella. This usually results in elicitation of a plant host defense response. However, rhizobia can be released into the cells of the developing nodule meristem, and a normally indeterminate nodule is formed (44).

The route of infection is characteristic for the host, because the same bacteria can penetrate different host species by either crack entry, infection through intact epidermis cells, or root hair infection threads. A given legume is infected by the same type of mechanism regardless of the infecting strain (40, 206). Similarly, the structural and developmental characteristics of an efficient nodule are specified by the plant and not by the rhizobial strain, indicating that the host possesses the genetic information for symbiotic infection and nodulation and that the role of the bacteria is to switch on this plant developmental program (for a review, see reference 53).

Rhizobial Genes Controlling Infection, Nodulation, and Host Range: an Overview

The *Rhizobium* genes essential for infection and nodule formation can be divided into two classes. One class includes several sets of genes involved in the formation of the bacterial cell surface, such as genes determining the synthesis of exopolysaccharides (*exo* genes), lipopolysaccharides (*lps* genes), capsular polysaccharides or K antigens, and β -1,2-glucans (*ndv* genes) (21, 95, 148, 183, 203). Mutations in these genes disturb the infection process to various degrees, such as the inability to elicit the formation of infection threads, resulting in the formation of nonfixing empty nodules ($\text{Nod}^+ \text{Fix}^-$ phenotype) (5, 149, 183, 201). A possible role of *exo* and *lps* genes in the determination of host specificity has been suggested, but no clear genetic evidence has yet been given that *Rhizobium* surface components are major determinants of host range specificity (64, 95, 183, 202).

The second class consists of the nodulation (*nod* or *nol*) genes. Inactivation of the nodulation genes can result in various in planta phenotypes, such as the absence of nodulation (Nod^-), a delayed but effective nodulation ($\text{Nod}^d \text{Fix}^+$), or changes in the host range. Some of the *nod* genes appear to be interchangeable for nodulation function between different species and biovars and are therefore designated as common *nod* genes (137). On the other hand, some *nod* genes are involved in the nodulation of a particular host and are hence called host-specific *nod* (*hsn*) genes (137). In most *Rhizobium* species studied to date, the *nod* genes reside on large symbiotic plasmids (pSym) that also carry the *nif* and *fix* nitrogen-fixing genes (166). In *Rhizobium loti* and *Bradyrhizobium* and *Azorhizobium* spp., the symbiosis-related genes are localized on the chromosome (3, 34, 85). Most *Rhizobium nod* genes are not expressed in cultured cells but are induced in the presence of the plant (for reviews, see references 75 and 136). This induction is caused mostly by flavonoids secreted by the plant (for reviews, see references 187 and 188) and also requires the participation of the transcriptional-activator protein NodD.

During the last few years, it has become clear that a major function of the *nod* genes is to ensure signal exchange between the two symbiotic partners (Fig. 1). In the first step, flavonoids excreted by the plant induce, in conjunction with the NodD protein, the transcription of bacterial *nod* genes (75, 221). In the second step, the bacterium, by means of the structural *nod* genes, produces lipooligosaccharide signals (Nod factors) (52a, 53, 240) that induce various root responses (245, 263). Mechanisms underlying host specificity depend on both the regulatory and the structural *nod* genes and will be discussed in further detail below.

STRUCTURAL *nod* GENES AS HOST RANGE DETERMINANTS

Common *nod* Genes

The structural *nod* genes are classified into two groups, the common and host-specific *nod* genes (137). The common *nodABC* genes have been found in all *Azorhizobium*, *Rhizobium*, and *Bradyrhizobium* isolates studied so far (85, 166, 251). These genes have been called common *nod* genes because they are structurally conserved and functionally interchangeable between *Rhizobium*, *Azorhizobium*, and *Bradyrhizobium* species without altering the host range (166). In most species, the *nodABC* genes are part of a single operon (Fig. 2). However, in *R. etli*, *nodA* is separated by approximately 20 kb from the *nodBC* genes (274). Inactivation of the *nodABC* genes abol-




FIG. 1. Schematic representation of the interaction between *Rhizobium* species and legume roots. Plant-secreted flavonoids, in conjunction with the bacterial NodD protein, regulate the transcription of bacterial *nod* genes, here exemplified by the *nodABCDEF* genes of *R. leguminosarum*. The bacterial *nod* gene products are involved in the synthesis of Nod factors. These Nod factors are able to induce the critical steps leading to nodule formation: (a) root hair deformation; (b) cortical cell division; (c) at least some steps of the infection process; and (d) increased flavonoid production. Reprinted with permission of the publisher from reference 78.

ishes the ability to elicit any symbiotic reaction in the plant, including root hair curling (Hac^-), infection thread formation (Inf^-), cortical cell divisions, and nodule formation (Nod^-), regardless of the host, the mode of infection, the type of nodule development, and the nodule location (159, 166).

The *nodIJ* genes are present in *R. leguminosarum* bv. *viciae* and bv. *trifolii*, *B. japonicum*, *R. etli*, and *A. caulinodans* (70, 80, 90, 257, 275), and partial sequence data suggest that they are present in *R. meliloti* (123). They reside downstream of *nodC* and seem to be part of the same operon as *nodC*. Mutations in *nodIJ* result in a delay of nodulation in *R. leguminosarum* and *A. caulinodans* (70, 80) but have no detectable effect in *B. japonicum* (90). It has been proposed that NodI and NodJ proteins belong to a bacterial inner membrane transport system of small molecules (275).

Host-Specific *nod* Genes

Other nodulation genes have been identified that are not functionally or structurally conserved among rhizobia. These host-specific *nod* (*hsn*) genes are necessary for the nodulation of a particular host plant (137). In most cases, mutations cannot be fully complemented by the introduction of the corresponding genes from other rhizobia. Mostly mutations result in alteration or extension of the host range. *R. meliloti nodH* mutants result in a change of host range; they infect and nodulate vetch but fail to nodulate their normal host, alfalfa. The *nodQ* mutants, on the other hand, are able to infect both alfalfa and vetch (71, 118). In *R. leguminosarum* bv. *viciae* and bv. *trifolii*, the *nodE* product is the main factor that distinguishes the host range of nodulation. In *R. leguminosarum* bv. *trifolii*, in contrast to the wild type, *nodEF* mutants nodulate white and red clover poorly but have acquired the ability to infect and nodulate peas. When these *nodEF* mutants of *R. leguminosarum* bv. *trifolii* harbor the *nodE* gene of *R. legu-*




FIG. 2. Genetic organization of *nod* genes in *R. leguminosarum* bv. *viciae* (a), *R. leguminosarum* bv. *trifolii* (b), *R. meliloti* (c), and *B. japonicum* (d). The genes are presented as arrows which point in the direction of their transcription. Common *nod* genes are indicated by solid arrows, and host-specific *nod* genes are indicated by hatched arrows. The *nodD* genes are indicated by white arrows. *nol* genes, unknown open reading frames, and other *nod* loci are indicated by stippled arrows. The *nodX* gene of *R. leguminosarum* bv. *viciae* TOM is not indicated but is located downstream from *nodJ* and presumably in the same operon (41). Black triangles indicate the position of the *nod* boxes. Updated from reference 217.

minosarum bv. *viciae*, they have an extended host range to *Vicia* and *Lathyrus* species (246). In *Rhizobium* sp. strain NGR234, mutation of *nodS* causes a Nod⁻ phenotype on *Leucaena leucocephala* but gives a normal phenotype on *Macropodium atropurpureum* (152). In a similar way, the *nodS* mutations in *R. tropici* CIAT899 and *A. caulinodans* cause a loss of nodulation of *L. leucocephala* (280). Waelkens et al. (280) also demonstrated that the *nodS* mutation of *Rhizobium* sp. strain NGR234, *R. tropici* CIAT899, and *A. caulinodans* exhibits a Nod⁻ phenotype on *Phaseolus vulgaris*. The *nodSU* genes have also been identified in *B. japonicum* and *R. fredii*, but no phenotype has been described for these genes (90, 143). Krishnan et al. (143) postulated that poor expression of the *R. fredii* *nodS* gene is the cause of the lack of *Leucaena* nodulation. Several loci involved in host-specific nodulation in *B. japonicum* have been identified. *nodVW* mutants of *B. japonicum* have lost the ability to nodulate mung bean (*Vigna radiata*), cowpea (*V. unguiculata*), and siratro (*Macropodium atropurpureum*), but the nodulation of soybean (*Glycine max*) is only marginally delayed (89). The *nodZ* gene of *B. japonicum* exhibits a host-specific characteristic at both species and cultivar levels, since *nodZ* mutants fail to nodulate siratro and shows an altered nodulation ability on a few varieties of soybean (182, 252).

Within the classic cross-inoculation groups of the *Rhizobium*-legume symbiosis, examples have been seen of more precise host-specific interaction at the plant cultivar level. Bacterial genes appear to control this cultivar specificity by acting as positive or negative traits. The first report on such a positive-acting gene was for the *R. leguminosarum* bv. *viciae*-pea symbiosis, in which a single gene, *nodX* from strain TOM, was identified as being responsible for overcoming the nodulation resistance of the cultivar Afghanistan peas (41). A similar case is the *nolA* gene identified in *B. japonicum* USDA110. The presence of *nolA* allows serocluster 123 isolates to nodulate soybean plants having USDA123-restricting genotypes. The nodules formed, however, were ineffective for symbiotic nitrogen fixation (211).

An example of a nodulation gene that acts negatively on the nodulation ability of a certain cultivar is *nodM*, present in *R.*

leguminosarum bv. *trifolii* TA1. The ability of strain TA1 to nodulate the subterranean clover cultivar Woogenellup is cold sensitive; at 28°C strain TA1 forms an effective symbiosis with this cultivar, but at 22°C little or no sign of nodulation is apparent (84). However, inactivation of *nodM* restores normal nodulation of cv. Woogenellup by strain TA1 (154). On the other hand, the introduction of *nodM* from the related *R. leguminosarum* bv. *trifolii* ANU843, which is able to fully nodulate cv. Woogenellup, again suppresses the nodulation of cv. Woogenellup by TA1, which indicates that the *nodM* genes of the two strains are interchangeable (154). In addition, the *nodT* gene from *R. leguminosarum* bv. *trifolii* ANU843 can act as a dominant suppressor of the negatively acting *nodM* and allows *R. leguminosarum* bv. *trifolii* TA1, which lacks a *nodT* gene, to nodulate cv. Woogenellup (155). In strain ANU843, however, there is no discernible phenotype for NodT recognized in nodulation of cv. Woogenellup (257).

R. fredii USDA257, a soybean symbiont, is able to form nodules on primitive soybean varieties, such as Peking, but fails to nodulate several agronomically advanced varieties, like McCall (107, 130). Recently, several negative-acting genes present in strain USDA257 have been identified. For instance, a Tn5 insertion in *nolC*, a chromosomal locus, enables strain USDA257 to form Fix⁻ nodules on soybean cv. McCall, but it also exhibits abnormalities on cv. Peking and expresses other pleiotropic effects (145). In contrast, the second group of Tn5 mutants, inactivating one of the genes *nolB*, *nolT*, *nolU*, *nolV*, *nolW* or *nolX*, all located on the pSym plasmid, enables *R. fredii* USDA257 to form fully fixing nodules on cv. McCall (170). Recently, Krishnan and Pueppke (146) discovered that *R. fredii* USDA257 is able to form nodules on *Erythrina costaricensis* but not on six other *Erythrina* species. Inactivation of *nolBU* or *nolC* broadens the symbiosis to include other *Erythrina* species (146).

In several cases, *nod* genes have been identified in mutants that exhibit no apparent defects in nodulation. For example, mutations of *nodT* in *R. leguminosarum* bv. *viciae* and bv. *trifolii* ANU843 (257) and of *nodSU*, *nodY*, and *nolYZ* in *B. japonicum* (90, 93) are not associated with an apparent altered

R. leg	AT	ATC	CATTCCATA	GAT	GATTGCC	ATC	CAAACAATC	AAT	TTTACCA	ATC	TTTCGGATC	ACT	TATAGAA
R. tri	CG	ATC	CACGCTGTA	GAT	GATTGCC	ATC	CAAACAATC	AAT	TTTACCA	ATC	TTTCGGAGT	GCT	TATTAGA
R. mel	GC	ATC	CATATCGCA	GAT	GATCGTT	ATC	CAAACAATC	AAT	TTTACCA	ATC	TTGCAGAGT	CCT	ATTAGAG
R. etli	GC	ATC	CATTGGACG	GAT	GAGTCC	ATC	CAAACAAT	GAT	TTTACCA	GCT	TATACCACT	GCC	ATTAGAA
B. jap	CT	ATC	CATCGTGTG	GAT	GTGTCT	ATC	GAAACAATC	GAT	TTTACCA	AAC	TGGGGGAGG	TTG	GATAGCA
A. caul	CC	ATC	GATCACGTG	GAT	TGGCTGT	ATT	CGGTAATTG	GAA	TTGACCG	GTA	GAATGATGG	TGC	ATAATTC
Consensus	Y	ATC	CAY...YR	GAT	G....Y	ATC	.AAACAATC	RAT	TTTACCA	ATC	Y		

FIG. 3. Sequence comparison of six *nod* boxes upstream from the common *nod* gene operons in *R. leguminosarum* bv. viciae (R.leg) (231), *R. leguminosarum* bv. trifolii (R.tri) (224), *R. meliloti* (R.mel) (209), *R. etli* (274), *B. japonicum* (B.jap) (182), and *A. caulinodans* (A.caul) (88). The consensus sequence was defined by Spink et al. (243). The ATC_NGAT repeats, or related motifs, are indicated in boldface type. Y, pyrimidine; R, purine; •, arbitrary base. The pairs of tandem arrows indicate the 9-bp repeats in the alternative *nod* box proposed by Wang and Stacey (282). The solid lines indicate the segments shown by interference footprinting to be critical for DNA binding according to Fisher and Long (76).

nodulation phenotype. These results are unexplained, but it is possible that the plants used in the assay are relatively insensitive to certain types of symbiotic effects. Alternatively, it is possible that the mutations are critical for the nodulation of some cultivars of plants not yet used in the assays.

REGULATORY *nod* GENES AS HOST RANGE DETERMINANTS

nodD Gene

The expression of the *Rhizobium* structural *nod* genes requires the presence of a plant signal, generally flavonoids, and the presence of the regulatory protein NodD. The NodD protein binds to conserved DNA sequences upstream of the inducible *nod* operons, called *nod* boxes. In the presence of plant signals, the NodD protein acts as a transcriptional activator (74, 209). This NodD regulatory system is present in all *Rhizobium*, *Bradyrhizobium*, and *Azorhizobium* strains studied so far. However, there are variations between species in the number of *nodD* copies present, and as many as five copies have been observed in some species (87, 159, 271a).

NodD as a member of the LysR family of transcriptional activators. The different *nodD* genes are conserved at the nucleotide sequence level, and the corresponding proteins have been classified as members of the LysR family of transcriptional activators (106). This group of regulator proteins shares several common features, as described by Schlaman et al. (222). These proteins all require an inducing compound for activation and all possess a putative helix-turn-helix motif in the amino region, characteristic for the DNA-binding ability. The NodD protein in *R. leguminosarum* bv. viciae and *R. meliloti* has been found to bind to the *nod* box, even in the absence of an inducer (74, 113). However, in *R. meliloti* AK41 and *Azorhizobium* species binding of NodD to the *nod* promoter is enhanced in the presence of its inducer (88, 139). The *nod* box was originally defined in *R. meliloti* as a 47-bp consensus sequence required for *nod* gene induction (209). In the genetically distant *B. japonicum* and *A. caulinodans*, less highly conserved *nod* boxes have been identified and shorter consensus sequences have been proposed (88, 282). A comparison of various divergent *nod* boxes revealed the presence of two inverted repeats ATC_NGAT (Fig. 3), which have a repeat structural feature typical for DNA targets that are symmetrically bound by protein dimers or tetramers (184). The presence in all *nod* boxes of such structures favors the hypothesis that NodD binds to the *nod* box as multimers (88). This model, however, is in disagreement with the model proposed by Fisher and Long (76). They provide strong arguments, based on interference footprints and DNA-phasing analysis, for two separable binding sites for NodD on one face of the DNA helix, in a region corresponding to the original consensus *nod* box (Fig. 3). This region does not contain completely the two paired

repeats ATC_NGAT, which lie on opposite faces of the DNA helix, as indicated in Fig. 3.

The specificity of each different NodD occurs at the level of amino acid sequence. Several hybrid NodD proteins have been constructed and exhibit the flavonoid specificity of the NodD product constituting their C-terminal end (116, 248). This indicates that the signal specificity is located primarily at the C-terminal part of the protein, which is less highly conserved than the N-terminal part. In addition, the carboxyl part of the protein exhibits some resemblance to animal steroid receptors, which are also known to interact with some flavonoid ligands (100). However, some of the mutations that modify the flavonoid specificity map in the N-terminal region (169, 248). Thus, the flavonoid specificity depends on the overall tertiary structure of NodD protein.

NodD as a membrane protein. In *R. leguminosarum* bv. viciae, the NodD protein is localized in the bacterial cytoplasmic membrane, presumably inserted only in the cytoplasmic monolayer (222). In *R. meliloti*, the NodD protein has been localized in the cytosol and migrates toward the cytoplasmic membrane when appropriate flavonoids are added (139). Also, flavonoids have been reported to accumulate in the cytoplasmic membrane (119, 196). Recently, Hubac et al. (119) have shown that the absorption of luteolin by *R. meliloti* in the inner and outer membranes involves the NodD proteins. These observations support the suggestion that the site of interaction between NodD and the flavonoid is likely to occur in the inner membrane (221). So far, there is no direct evidence that flavonoid inducers bind directly to NodD proteins. However, genetic evidence is in line with this hypothesis: (i) point mutations in *nodD* may change the sensitivity to additional inducers (25, 26, 169, 248); and (ii) the transfer of a *nodD* gene to a foreign *Rhizobium* strain is linked to the transfer of sensitivity to a given set of flavonoids (116, 247). Since flavonoids are required for activation of the NodD protein, they presumably induce a conformational change in the protein. This notion is supported by the fact that it is possible to construct mutants and hybrid *nodD* alleles which can activate transcription in the absence of flavonoids (25, 244).

In *R. leguminosarum* bv. viciae and *R. meliloti*, the soluble form of NodD binds to *nod* box sequences even in the absence of inducers, although in *R. leguminosarum* bv. viciae a minor fraction of cytoplasmic membrane-located NodD is able to bind to the *nod* box as well (74, 217). Using migration retardation assay with the flavonoid-independent *nodD3* allele of *R. meliloti* Fisher and Long (76) have shown that NodD3 induces or stabilizes a bend in the *nod* box upon binding and that the bend center lies between the two NodD-binding sites. It is possible that the specific bend induced by NodD3 is immediately present in an "active" mode and that this protein bends the *nod* box more or less than the flavonoid-dependent NodD proteins do (76). In *R. meliloti*, a chaperone-like protein showing homology to GroEL of *Escherichia coli* is necessary for the

TABLE 2. Effects of flavonoids, in conjunction with various NodD proteins, on *nod* gene expression^a

Compound	Substitution position							Activity of NodD protein ^b							
	3	4	5	7	3'	4'	5'	RtrD1	Rt	Rl	RmD1	RmD2	NGR234	Bj	RpD2
Flavones															
Luteolin			OH	OH	OH	OH		+	++	++	++	—	++	—	+
Apigenin			OH	OH	OH			++	++	++	+	—	++	+	++
— ^c				OH	OH	OH		ND	ND	++	++	—	ND	ND	ND
Chrysoeriol			OH	OH	OCH ₃	OH		ND	ND	ND	++	—	ND	ND	+
—					OH	OH		ND	ND	ND	—	—	ND	ND	ND
—				OH		OH		++	++	ND	+	ND	++	+	ND
Chrysin			OH	OH				++	+	—	—	—	++	—	+
Galangin	OH		OH	OH				—	—	—	ND	ND	ND	ND	+
Flavonols															
Myricetin	OH		OH	OH	OH	OH	OH	—	—	—	—	—	+	ND	ND
Quercetin	OH		OH	OH	OH	OH		—	—	—	—	—	++	—	ND
Kaempferol	OH		OH	OH		OH		—	—	—	—	—	++	+	+
Flavanones															
Eriodictyol			OH	OH	OH	OH		—	+	++	+	—	ND	—	+
Naringenin			OH	OH		OH		++	++	++	—	—	++	—	++
Hesperitin			OH	OH	OH	OCH ₃		—	—	++	—	—	++	ND	+
Isoflavones															
Genistein			OH	OH		OH		—	—	—	—	—	++	++	++
Daidzein				OH		OH		ND	—	—	—	—	++	++	ND
Chalcone															
4,4'-Dihydroxy-2'-methoxychalcone		OH				OH		ND	ND	ND	++	++	ND	ND	ND

^a Data from references 43, 53, and 270.^b The *nod* inducing activities of various flavonoids were recorded for rhizobial strains harboring the *nodD* genes of *R. tropici* NodD1 (RtrD1); *R. leguminosarum* bv. trifolii (Rt), *R. leguminosarum* bv. viciae (RI); *R. meliloti* NodD1 (RmD1) and NodD2 (RmD2); *B. japonicum* (Bj) and *R. leguminosarum* bv. phaseoli (Rp). The different activities are related to the maximum inducing activity recorded for these strains. ++, 50 to 100%; +, 10 to 50%; —, 0 to 10% of the maximum induction. ND, not detected.^c —, compounds for which no commercial name is available yet.

transcriptional activity of NodD3 (160). It is conceivable that this protein is necessary for the translocation of NodD from the cytoplasmic membrane to keep it in a proper, soluble conformation (221).

Plant signals activating NodD. The activation of *nod* expression by NodD proteins requires, with some exceptions, the presence of compounds in plant exudates (122, 175, 208). They were first identified as flavonoids or related compounds derived from plant phenylpropanoid metabolism, which is also known to provide molecules involved in plant defense (73, 185, 197). Flavonoids are released as aglycones or glycosidic conjugates. The latter are less active but have a higher solubility in water, and they can be converted to active forms by bacterial glycosidases (104, 120, 168). Recently, Phillips et al. (189) identified nonflavonoids, namely, the two betaines trigonelline and stachydrine, as being two major compounds from alfalfa seed exudates activating the NodD2 protein in *R. meliloti*. Betaines are chemically quite different from flavonoids, have different physicochemical properties (for example, water solubility) and are synthesized by different metabolic pathways in the host plants. It is interesting that these two types of molecules (flavonoids and betaines), which are chemically so different, are able to cause transcriptional activation of the isoforms of the same protein.

The nature and amounts of the compounds exuded depend on the plant and its stage of development. For alfalfa, soybean, and bean, the spectrum of flavonoids present in seed exudates is different from that present in root exudates (94, 103, 120, 121). Inoculation with an infective rhizobial symbiont causes a change in the internal flavonoid pool of the root. This second wave of flavonoids leads to an increased *nod* gene-inducing activity, by as much as 10-fold, as detected for white clover (207), vetch (268), soybean (223), bean (39), *Lotus* species

(35), and alfalfa (38). In several cases, flavonoids without inducing properties have been shown to inhibit *nod* gene activation by effective inducers (54, 73, 101, 141, 186). The anti-inducers usually have similar structures to those of the inducers, and inhibition can be overcome by increasing the concentration of the inducers (186). Also, synergistic interactions have been observed and can be explained if one considers that NodD acts as a multimer (103, 105).

NodD as a determinant of host range. It is apparent that the spectrum of flavonoid specificity of the endogenous NodD protein correlates with the broadness of the host range. NodD proteins from narrow-host-range rhizobia, like *R. meliloti*, *R. leguminosarum* bv. viciae, and *R. leguminosarum* bv. trifolii, respond to few flavonoids, while NodD from the broad-host-range *Rhizobium* NGR234 has a larger spectrum of inducing compounds, including even the monocyclic aromatic compounds vanillin and isovanillin (101, 151) (Table 2).

Several lines of genetic evidence have established that *nodD* genes, because of the flavonoid specificity, are determinants of host specificity. (i) Some mutations in *nodD* cannot be complemented by a *nodD* from other *Rhizobium* species (116, 247). For example, the *R. meliloti nodD1* gene cannot complement the *nodD* mutation of NGR234 for nodulation of siratro. This lack of complementation is due to the inability of *R. meliloti nodD1* to respond to the signals exuded by siratro. (ii) Transfer of the *nodD* gene alters the host range. The transfer of the *nodD1* gene from *Rhizobium* sp. strain NGR234 into *R. meliloti* results in the transfer of the ability to nodulate siratro (116). (iii) Some point mutations in the *nodD* gene of *R. trifolii* can extend the host range to the nonlegume *Parasponia* species (169). (iv) A *nodD* hybrid gene, constructed in vitro from *R. meliloti* and *R. trifolii nodD* genes, extends the host range to include tropical legumes (244). These experiments indicate

that different NodD proteins have different inducer specificities and, as a consequence, influence the host specificity of nodulation (116, 247).

NodD regulation. The *nodD* gene exists as a single gene in *R. leguminosarum* bv. *viciae* and bv. *trifolii* (59, 122) and probably also in *Azorhizobium* species (87). Inactivation of the *nodD* gene consequently confers a Nod⁻ phenotype. Other rhizobia possess multiple copies of the *nodD* gene: *R. fredii* USDA 191, *Rhizobium* sp. strain NGR234, and *B. japonicum* harbor two copies, of which NodD1 is the activator of *nod* genes whereas no such a function could be assigned to NodD2 (3, 23, 91). *R. meliloti* and *R. leguminosarum* bv. *phaseoli* possess three *nodD* genes (42, 92, 115). Mutation in one of the three *nodD* homologs of *R. meliloti* reduces nodulation in a host-dependent manner (99, 114). Up to five copies of the *nodD* gene have been identified in *R. tropici* CIAT899 (271a).

The significance, in terms of control of specificity, of these *nodD* reiterations is not clear, because there is no correlation between the number of *nodD* genes and the broadness of the host range: the narrow-host-range *R. leguminosarum* bv. *phaseoli* has three copies (42), whereas the broad-host-range *Rhizobium* sp. strain NGR234 possesses two *nodD* copies, of which only one seems to be involved in *nod* gene induction (23). It is possible that NodD plays other roles besides induction of *nod* genes. This can be explored only with strains having multiple *nodD* copies by analysis of mutants at various levels of bacterium-plant interactions.

The various *nodD* genes also differ in their regulation. The single *nodD* gene negatively regulates its own transcription in *R. leguminosarum* bv. *trifolii* and bv. *viciae* (208, 243). In *B. japonicum*, *R. leguminosarum* bv. *phaseoli*, and *R. fredii*, the *nodD1* gene is preceded by a *nod* box sequence (3, 43, 282). For the first two species, the *nodD1* transcription level is enhanced in the presence of NodD1 protein and certain flavonoids, independently of other *nod* genes (3, 43).

In *R. meliloti*, the expression of *nodD3* and *syrM*, another member of the LysR family, is strongly intertwined in a complex way: the *syrM* gene product activates the expression of *nodD3*, which in turn activates the expression of *syrM* (138, 210, 259). The two genes constitute a self-amplifying positive regulatory circuit. Whereas NodD1 and NodD2 are specifically activated by plant compounds, plant inducers have little effect on the *syrM-nodD3* interaction. In the free-living state, *R. meliloti* strains with *syrM* and *nodD3* present in a single copy show no detectable expression of either gene (259). However, when both are carried on a multicopy plasmid, *SyrM* and *NodD3* are present at sufficiently high concentrations to induce high levels of *nodABC* expression in the absence of an inducer (176, 259). This suggests that one or both genes might be repressed in the free-living state (259). To a lesser extent, the *nodD2* gene influences *syrM* expression (138) and *nodD1* influences *nodD3* expression (162). Interestingly, *SyrM* also regulates the synthesis of the *exo* genes, indicating that *SyrM* could coordinately regulate the metabolism of exopolysaccharide and of the Nod factors, both of which are involved in the infection process (12, 210). Recently, *syrM* has also been identified in *Rhizobium* sp. strain NGR234 and *R. leguminosarum* bv. *phaseoli* (173, 198).

Another role of NodD seems to be the regulation of *nod* genes as a function of the concentration of combined nitrogen. *R. meliloti* NodD3 and *B. japonicum* NodD1 are involved in the repression of the *nod* regulon in the presence of an excess of combined nitrogen (66, 281). In *R. meliloti* two NtrC-binding sites were found upstream of *nodD3* (135), whereas in *B. japonicum* neither *NifA* nor *NtrC* appears to be involved.

A repressor of *nod* gene expression, *NolR*, has been identified in several strains of *R. meliloti*, but no *NolR* function could

be found in the well-studied strain *R. meliloti* 1021. *NolR* binds to the promoter region of *nodD1* and *nodD2* and not to any of the inducible *nod* promoters. Upon binding, it controls negatively the expression of these two regulatory genes and consequently of the *nod* regulon (139, 140). A mutation in the repressor results in a delayed nodulation phenotype. Thus, in some strains, *nolR* controls the fine-tuning of *nod* gene expression (139). In *R. meliloti* 1021, the lack of *NolR* function is shown to be due to a single insertional mutation in the C-terminal coding sequence which abolishes the DNA-binding activity (37). Sanjuan et al. (213) isolated, by hybridization to the *R. meliloti* *nolR* locus, a corresponding repressor locus in *B. japonicum*. Mutations in this loci result in an elevated level of *nod* gene induction. Hybridization studies suggest that *nolR* does not exist in *R. leguminosarum* bv. *viciae*. It was shown, however, that the chromosomal background in *R. leguminosarum* strongly influences the expression of *nod* genes, indicating that an additional regulator may act in some *R. leguminosarum* strains as well (220). In *A. caulinodans*, at least three other proteins, smaller than NodD, were found to bind to *nod* box DNA, but their function is unknown (88).


The inducible *nod* genes are not transcribed in the bacteroids, and thus their expression stops after the bacteria are released from the infection thread (218, 230). Apparently, in *R. leguminosarum* bv. *viciae* this is caused by ineffective binding of NodD to the *nod* box, because of either a conformational change of the protein or its presence in another complex (219). In the bacteroids of *R. leguminosarum* bv. *viciae*, the level of *nodD* expression is approximately 35% of that of the free-living cells, and this may be caused by a bacteroid-specific repressor protein (219). In *R. meliloti*, the *nodD3* gene is expressed in the meristematic region and the infection zone of the nodule, while *syrM* is expressed in the central tissue of the nodule (230). Swanson et al. (259) showed by in situ expression studies the interdependence of *nodD3* and *syrM* expression in the nodule: no *nodD3* expression occurred if *syrM* was mutated, and, conversely, no *syrM* expression was detected if *nodD3* was mutated.

Other *nod* Regulatory Genes

Other *nodD* regulatory genes, which are determinants of host specificity, have been found in *Bradyrhizobium* species. Mutants with mutations of *nodVW* have lost the ability to nodulate cowpea, mung bean, and siratro, but nodulation on soybean is only marginally delayed (89). The predicted amino acid sequence of NodVW suggests that these gene products are members of the family of two-component regulatory systems. This led to the hypothesis that NodV responds to an environmental stimulus and that after signal transduction, NodW may be required to positively regulate the transcription of one of several unknown genes involved in the nodulation process (89). Recently, Sanjuan et al. (214) found that *nodW*, in conjunction with *nodD1*, is essential for maximal induction of the common *nod* genes and *nodD1*. How the NodW protein is involved in the regulation is not known. The possibility that other, as yet unknown, genes are under control of NodW has not been ruled out. Also, the possibility that NodW is involved in modifying plant signals required for NodD activation of *nod* promoters should be considered (214). Grob et al. (96) reported the identification of a *B. japonicum* gene, *nwsB*, a *nodW* suppressor, which when overexpressed from a strong promoter is able to suppress the Nod⁻ phenotype from the *nodW* mutant. The *nwsB* gene is preceded by a long open reading frame, *nwsA*. The protein products of the *nwsAB* gene pair also appear to be two-component regulators and are most similar to

TABLE 3. Structures of lipooligosaccharides produced by different *Rhizobium* strains^a

Strain	<i>n</i>	Q	R ₁	R ₂	R ₃	R _{4,5}	Reference
<i>R. leguminosarum</i> bv viciae RBL5560	2, 3	C _{18Δ2 4 6 11} C _{18Δ11}	CH ₃ CO	H	H	H	245
<i>R. meliloti</i> AK41	1, 2, 3	C _{16Δ2 9} C _{16Δ2 4 9}	H CH ₃ CO	SO ₃ H	H	H	226
<i>R. meliloti</i> 2011	2, 3	C _{16Δ9} C _{16Δ2 9} C _{16Δ2 4 9} (Ω-1)-OH C ₁₈₋₂₆	H CH ₃ CO	SO ₃ H	H	H	51, 52, 150
<i>R. tropici</i> CFN299	3	C _{18Δ11}	H	SO ₃ H	CH ₃	H	191
<i>B. japonicum</i> USDA110	3	C _{18Δ9}	H	2-O-Me-Fuc	H	H	212
<i>B. japonicum</i> USDA135	3	C _{18Δ9}	H	2-O-Me-Fuc	H	H	31
<i>B. japonicum</i> USDA61	2, 3	C ₁₆ C _{18Δ9}	CH ₃ CO H CH ₃ CO	Fuc 2-O-Me-Fuc	H CH ₃	H	31
<i>R. fredii</i> USDA257	1, 2, 3	C _{18Δ11}	H	Fuc 2-O-Me-Fuc 2-O-Me-Fuc	H	H	14
<i>Rhizobium</i> sp. strain NGR234	3	C _{18Δ11} C ₁₆	H	2-O-Me-Fuc 2-O-Me, 4-O-SO ₃ H-Fuc 2-O-Me, 3-O-CO-CH ₃ -Fuc	H	NH ₂ CO	192
<i>A. caulinodans</i> ORS571	2, 3	C _{18Δ11} C ₁₈	H NH ₂ CO	H D-Arabinosyl	CH ₃	H	172



^a The chitin oligomer and the acyl moiety (Q) are present in all Nod metabolites. The number (*n*) of *N*-acetylglucosamine residues can vary. Q can vary in length and in the number of unsaturated bonds. Several different substitutions to the sugar backbone occur (R₁ to R₅), which are listed in the table. Me-Fuc, methylfucose.

the protein products of the *B. japonicum* gene pair *nodVW* (97). Cross talk occurs between NwsAB and NodVW, since the activity of NwsB depends on either the NodV or the NwsA sensor kinase. The physiological importance of cross talk between NodVW and NwsAB in *B. japonicum* remains unclear (97).

Another gene of *B. japonicum*, *nolA*, plays a role in the nodulation of soybean genotypes that are restricted in nodulation by members of *B. japonicum* serocluster 123 (211). The N terminus of the predicted gene product of *nolA* has strong similarity with the N terminus of MerR, the regulator of mercury resistance genes. Although no *nod* box sequence could be found upstream of *nolA*, the expression of *nolA* is moderately induced by soybean seed extracts and the isoflavone genistein (211). The *nodZ* gene of *B. japonicum* exhibits a host-specific function for the nodulation of siratro and a few varieties of soybean (182, 252). Functional analysis showed that *nodZ* is involved in the production of the Nod factor but is not regulated by NodD. Surprisingly, it shows an elevated expression in the bacteroids, although NifA is not required for *nodZ* expression and expression is not influenced by oxygen concentration (252).

Several *nod* genes of *R. fredii* are not preceded by a *nod* box (20, 144, 170). *nolC*, a chromosomal locus of *R. fredii* USDA257, was first identified as a negative regulator of cultivar-specific nodulation of soybean cv. McCall (144). However,

further investigation revealed that *nolC* is also detrimental to nodulation of other soybean plants, like cv. Peking, has other pleiotropic effects (145), and acts as a negative regulator for specific nodulation of several *Erythrina* species (146). *nolC* has strong sequence homology to *dnaJ*, which encodes a heat shock protein in *Escherichia coli* and contains a regulation function (144). *nolC* is constitutively expressed, and in planta histological studies confirm that its expression is not temporally regulated but, rather, is sustained throughout nodule development, including the preinfection stage (145). A second gene cluster, consisting of *nolBTUV*, *nolW*, and *nolX*, which is present on the pSym plasmid, is involved in the cultivar-specific nodulation of soybean cv. McCall (170). Insertion mutations of each gene extend the host range to soybean cv. McCall without exerting any effect on other host plants. Also, the inactivation of *nolBU* broadens the host range of *R. fredii* USDA257 to include several new *Erythrina* species (146). Expression of *nolBTU* and *nolX* is induced as much as 30-fold by flavonoid signal molecules, even though these genes lack *nod* box promoters. In planta analysis verified that these genes are expressed continuously from preinfection to the stage of the functional nodule (170). Another *nod* box-independent *nod* gene has been identified in *R. fredii* USDA201, namely, *nolJ*, which requires a functional *nodDI* gene for induction (20). *nolJ* shares no sequence homology with any of the previously reported common or *hsn* nodulation genes.

Nod FACTORS

Rhizobia produce and secrete soluble factors in response to the inducing flavonoids from the plant. The *nod* genes are essential for this process (268, 269). Recently, the chemical structures of these nodulation factors, also called Nod factors, have been determined (14, 31, 150, 172, 191, 192, 212, 226, 245). Uniformly, they consist of an oligosaccharide backbone of β -1,4-linked *N*-acetyl-D-glucosamine varying in length from three to five sugar units. A structurally varied fatty acid group is attached to the nitrogen group of the nonreducing amino sugar moiety. The presence of other substitutions is dependent on the species and strain (Table 3).

Biological Activities of Nod Metabolites

Nod factors are considered to be the main *Rhizobium* nodulation signal molecules, since the purified molecules are able to induce many of the plant responses observed in early stages of symbiosis (245, 263). Lipooligosaccharides from several species of rhizobia are able to elicit root hair deformation and induce nodule primordia in a host-specific way, indistinguishable from the nodule meristem in the first stage of normal nodule organogenesis (105a, 150, 172, 200, 212, 245, 263). Normally, the formation of shepherd's crooks has not been detected by treatment with Nod factors. Only in the case of the very promiscuous *Macropitium atropurpureum* does the application of Nod factor provoke marked shepherd's crook-type curling of the root hair. This even occurs with Nod factors of *Rhizobium* strains which are not able to nodulate *M. atropurpureum*, although a higher concentration of Nod factor is needed (200). In *Medicago sativa* and *Glycine soja*, the nodule meristems are capable of developing further into full-grown nodules which have anatomical and histological features of genuine nodules induced by *Rhizobium meliloti* and *Bradyrhizobium japonicum* (254, 263). Recently, nodule-like structures following treatment with Nod factors have been observed with soybean, siratro, bean, and *Sesbania* species (165, 172, 200). Purified Nod factors are not able to promote the formation of genuine infection threads, but certain effects of the lipooligosaccharides have been observed which suggest that they are involved in the infection process. The formation of pre-infection thread structures is observed after treatment of *Vicia* roots with the lipooligosaccharides of *R. leguminosarum* bv. *viciae* (267). Another indication comes from the observation that the transcription of some early nodulins, which are specifically expressed in the infection thread, is also induced with isolated Nod metabolites (117, 128, 177, 215).

The lipooligosaccharides of *R. leguminosarum* bv. *viciae* are able to elicit the production and secretion of additional flavonoids in the roots of *Vicia sativa* (195). These additional flavonoids are efficient inducers of the NodD-dependent *nod* gene expression and are also found by treatment of plants with infective rhizobia (39, 268). In a similar case, the response of soybean roots to either *B. japonicum* inoculation or treatment with Nod factors of *B. japonicum* or *Rhizobium* sp. strain NGR234 resulted in the accumulation of flavonoid molecules in soybean root exudates (223).

It has been shown that application of cytokinin and auxin transport inhibitors induces the formation of pseudonodules similar to the formation induced by treatment of Nod factors (36, 109, 200). Since Nod factors share properties with plant hormones, they may be regarded as "hormone-like" molecules. However, Nod factors possess another essential characteristic, namely, the ability to curl root hairs. These two properties of Nod factors allowed the *nodABC* deletion mutant of *Rhizobium* sp. strain NGR234 and of *B. japonicum* to enter the roots

of *Macropitium atropurpureum* and *Glycine max*, respectively, in the presence of NGR234 Nod factor (199, 200). The Nod factor of NGR234 also allowed *R. fredii* USDA257 to enter and fix nitrogen on the nonhost *Calopogonium caeruleum* (199).

The activity of Nod factors can also be determined by the presence of plant enzymes involved in the metabolism of the Nod factor. Staehelin et al. (253) have shown that the Nod factors of *R. meliloti* are rapidly inactivated in the rhizosphere of alfalfa by the action of chitinases and that the rate of degradation depends on the structural modification of the Nod factor. The presence of the sulfate group on the O-6 position of the reducing end of the *R. meliloti* Nod factor strongly protects against degradation by purified plant chitinases and intact plant roots (253). The finding that the Nod factors are substrates for plant chitinases and the fact that the Nod factors are cleaved by chitinases at a different rate might be an important determinant in the host specificity (225).

Biochemical Function of the *nod* Gene Products

Knowledge of the biochemical function of several Nod gene products involved in the biosynthesis has been obtained in different ways: (i) comparison of the Nod factors of wild-type and mutant strains, (ii) prediction of their function by searches for structural homology with already known proteins, and (iii) physiological or biochemical study of the *nod* genes products. The information obtained by comparison of the deduced amino acid sequences with other proteins of known function is summarized in Table 4.

Common *nod* genes and the molecular backbone. It has been shown recently that the common *nodABC* genes are sufficient to synthesize the backbone of signal molecules consisting of *N*-acylated glucosamine oligosaccharides (245). Results obtained by Geremia et al. (83) from in vitro labelling studies with cell extracts of several *Azorhizobium* mutants and a construct that produces the NodC protein in *E. coli*, strongly suggest that NodC functions as an *N*-acetylglucosaminyltransferase involved in the synthesis of the chitin backbone. This conclusion is consistent with the results of homology studies, which indicated that positions of NodC are homologous with a domain of various chitin synthases and cellulose synthases (6, 24, 48). It is not yet clear how the exact number of monomers in the oligosaccharide backbone (tri-, tetra-, or pentamer) is controlled. This hypothesis of polymerization of *N*-acetylglucosamine subunits to form the oligosaccharide backbone is supported by the finding that the *nodM* protein has sequence homology with glucosamine synthase (encoded by *glmS*) (164) and therefore could have a function in the production of glucosamine, an obvious precursor of the synthesis of the oligosaccharide backbone. The predicted biochemical function of NodM is supported by the observation that the *nodM* gene was able to complement an *E. coli glmS* mutation (7). A mutation in *nodM* exhibits only a moderate delay in infection and nodulation (7, 255). This is consistent with the fact that a mutation in *nodM* results only in a quantitative decrease of the production of lipooligosaccharides (9, 243, 245). Leakiness might be due to the presence of a chromosomal counterpart of *nodM* (9, 164).

An important question is still which enzymes are responsible for the acylation of the chitin backbone. It is reasonable to assume that prior to the attachment of the correct fatty acyl chain to the nonreducing *N*-acetylglucosamine residue of the chitin backbone, the corresponding *N*-acetyl amino group must be deacetylated. Results from John et al. (125) indicate indeed that the NodB protein from *R. meliloti* deacetylates the nonreducing *N*-acetylglucosamine residue. Consequently, the last

TABLE 4. Some features of *nod* gene products

Nod protein	Species biovar ^a	Cellular location ^b	Sequence homology	Reference(s) ^c
NodA	Common	Cyt		6a, 126, 205
NodB	Common	Cyt	Deacetylase	6a, 125
NodC	Common	Inner mb	Chitin synthases	6, 13, 24, 48, 83
NodD	Common	Cyt mb	Transcription activator, LysR family	106, 222
NodE	Rl, Rt, Rm	Cyt mb	β -Ketoacyl synthase	17, 52, 246
NodF	Rl, Rt, Rm	Cyt	Acyl carrier protein	52, 81, 231
NodG	Rm		Alcohol dehydrogenase, β -ketoacyl reductase	49, 232
NodH	Rm		Sulfotransferase	204
NodIJ	Rl, Rt, Rm, Re, Ac, Bj, Bp	Cyt mb	Capsular polysaccharide secretion proteins	275
NolK	Bp			229
NodL	Rl, Rt, Rm, Bp	Cyt mb	Acetyltransferase	9, 18
NodM	Rl, Rt, Rm, Bp		D-Glucosamine synthase	8
NodN	Rl, Rt, Rm, Bp			8
NodO	Rl	Secreted	Hemolysin	258
NodP	Rm		ATP-sulfurylase	227
NodQ	Rm		ATP-sulfurylase and APS kinase	227
NodS	Re, Rtr, Rf, Bj, NGR, Ac		Methyltransferase (Ac)	80, 90, 152
NodT	Rl, Rt	Outer mb	Transit sequences	257
NodU	Re, Rtr, Rf, Bj, NGR, Ac			80, 90, 152
NodV	Bj	Cyt mb	Sensor, two-component regulatory family	89
NodW	Bj	Cyt	Regulator, two-component regulatory family	89
NodX	Rl*		Acidic exopolysaccharide encoded by <i>exoZ</i>	41, 72
NodY	Bj			182
NodZ	Bj			252
NolA	Bj		DNA-binding protein	211
NolB	Rf			170
NolC	Rf		Heat shock protein DnaJ	144
NolE	Rp	Secreted		42
NolFGHI	Rm			7
NolJ	Rf			20
NolK	Ac		Sugar epimerase	86
NolMNO	Bj			161
NolP	Rp			42
NolR	Rm		DNA-binding protein	140
NolT	Rf		<i>hrp</i> genes of pathogenic bacteria	170
NolUV NolW NolX	Rf			170
NolYZ	Bj			55

^a Nod proteins are present in *R. leguminosarum* bv. *viciae* (Rl), *R. leguminosarum* bv. *trifolii* (Rt), *R. leguminosarum* bv. *phaseoli* (Rp), *R. meliloti* (Rm), *R. tropici* (Rtr), *R. etli* (Re), *R. fredii* (Rf), *Rhizobium* sp. strain NGR234 (NGR), *B. japonicum* (Bj), *Bradyrhizobium* sp. strain *Parasponia* (Bp), *A. caulinodans* (Ac), and *R. leguminosarum* TOM (Rl*).

^b Cyt, cytoplasmic; mb, membrane.

^c Only references that describe the most recent data about subcellular location, biochemical function, or homology are given.

step in forming the lipooligosaccharide backbone structure is the N-acylation on the deacetylated nonreducing end, which has been shown to be carried out by NodA (6a, 205).

Nothing is known about the transport of lipooligosaccharides out of the cell. The NodI protein shares sequence homology with a large family of traffic ATPases and has therefore been proposed to be involved in the secretion of the lipooligosaccharides (58, 275). However, transposon mutations in the *nodI* gene did not have a quantitative effect on the amount of lipooligosaccharides released in the growth medium (242). Recent results, however, with a multicopy plasmid containing cloned *nod* genes indicate that *nodI* and *nodJ* could have an influence on the excretion of lipooligosaccharides (241).

Synthesis of specific side chains. Several *nod* genes that were shown to be involved in the determination of the host range specificity of nodulation also appear to be involved in the production of lipooligosaccharides. Most noteworthy are the genes *nodQ* and *nodH*, which are the major determinants of host specificity in *R. meliloti* (75, 226, 227), and the gene *nodE*, which was shown to be the major determinant of host specificity in *R. leguminosarum* bv. *viciae* and bv. *trifolii* (58, 245).

The sequence homologies of NodF with the acyl carrier proteins and of NodE with various β -ketoacyl synthases (17, 58) lead to the hypothesis that these gene products function in the synthesis of the highly unsaturated lipid moiety. This hypothesis was further supported by the observation that NodF contains a 4'-phosphopantetheine prosthetic group (81), which can function as a carrier for acyl chains during fatty acid biosynthesis or transfer. The *nodE* mutants of both *R. leguminosarum* bv. *viciae* and *R. meliloti* secrete Nod metabolites that are N-acylated by vaccenic acid (C_{18:1}), the most common fatty acid in gram-negative bacteria. In contrast, the wild-type strains also secrete Nod factors with a highly unsaturated fatty acid, C_{18:4} for *R. leguminosarum* bv. *viciae* and C_{16:2} or C_{16:3} for *R. meliloti* (52, 226, 245). This is explained in a simple model in which it is assumed that the NodE protein, by using regular fatty acid intermediates as substrates (which cannot be used by the normal household β -ketoacyl synthases), competes with normal fatty acid synthesis (82). In addition to Nod factors which are mono-N-acylated by unsaturated C₁₆ fatty acids, *R. meliloti* produces a minor group of Nod factors which contain a series of C₁₈ to C₂₆ (ω -1)-hydroxylated fatty acids. The syn-

thesis of these N-acyl substituents is under control of the regulatory genes *nodD3* and *sydM* but is not specified by *R. meliloti* *nodEF* genes (51, 52). In *R. meliloti*, the *nodG* gene, which is not found in *R. leguminosarum*, resides downstream from *nodEF* (49, 77, 118). NodG shares homology with β -ketoacyl reductases and several alcohol dehydrogenases and might be involved in the synthesis of the acyl moiety (48, 77, 118). However, *nodG* mutations in *R. meliloti* did not result in a detectable change in the *R. meliloti* Nod factor acyl chain, but it is still possible that another *Rhizobium* gene can substitute for NodG function in the mutated strain (52).

In *R. meliloti*, a mutation in the *nodH* gene results in the absence of the sulfonation of all lipooligosaccharides. Mutations in *nodP* and *nodQ* genes result in the production of a mixture of sulfated and nonsulfated Nod factors (204). This leaky phenotype seems to result from the presence of a second copy of the *nodPQ* genes on the second megaplasmid (228). *R. meliloti* strains carrying mutations in both *nodPQ* copies do not produce sulfated factors (204). In gram-negative bacteria, the first steps of sulfate metabolism involve the synthesis of activated forms of sulfate derived from ATP, first the formation of adenosine 5'-phosphosulfate (APS) by ATP sulfurylase and second the formation of 3'-phosphoadenosine 5'-phosphosulfate (PAPS) by APS kinase (142). The NodP protein and a region of NodQ share strong homology with CysD and CysN of *E. coli*, which are subunits of an ATP sulfurylase. Another region of *nodQ* is homologous to CysC, an APS kinase (228). In vitro studies have demonstrated that the NodP and NodQ proteins indeed have ATP sulfurylase and APS kinase activity (228), and their function is probably synthesis of PAPS. The NodH protein shares homology with sulfotransferases (204), which is consistent with its indispensable role in the addition of the sulfate moiety.

Mutations within *nodL* of *R. leguminosarum* strongly reduce nodulation on *Pisum*, *Lens*, and *Lathyrus* species but have only little effect on nodulation of *Vicia* species (256). *R. meliloti* *nodL* mutants exhibit only a slight delay of nodulation on alfalfa (53). NodL is homologous to bacterial acetyltransferases (8, 57), and in *R. leguminosarum* they specify the O acetylation on the C-6 of the nonreducing terminal glucosamine residue (18, 245). The same function is attributed to *R. meliloti* *nodL* (53).

Nod Metabolites: Variation on a Theme

Bradyrhizobium and *Azorhizobium* species are genetically quite distant from *Rhizobium* species (61, 124), but they were found to produce Nod metabolites belonging to the same family as those from *Rhizobium* (31, 172, 212).

Rhizobium sp. strain NGR234 produces a family of Nod metabolites which are also mono N-acylated pentamers of chitin with a variety of possible substitutions. The fact that NGR234 secretes a mixture of both sulfated and nonsulfated Nod metabolites may be part of the basis for its symbiotic promiscuity (192).

In *B. japonicum*, the Nod factors are also monoacylated chitin pentamers. However, the reducing end is substituted on C-6 with methylfucose (212). Results by Stacey et al. (252) show that the *nodZ* gene is responsible for the fucosylation of the core lipooligosaccharide nodulation signal. Since the Nod factors of *Rhizobium* sp. strain NGR234, also capable of soybean nodulation, have a 2-O-methylfucosyl (or fucose) residue (192), it suggests that this residue is critical for soybean nodulation. Indeed, the Nod factor isolated from a *B. japonicum* *nodZ* mutant is not able to induce cortical cell divisions on *Glycine soja* (252). However, the *nodZ* mutant is still able to

nodulate most soybean genotypes (182, 252). Therefore, it is possible that the 2-O-methylfucose substituent is vital to soybean nodulation but that other functions in *B. japonicum* are able to compensate for the loss of this substituent (252). Also, mutations in the *nolO* gene of *B. japonicum* result in an alteration in the profile of nodulation signals, produced upon induction with flavonoids. Moreover, *nolO* mutants of *B. japonicum* produce a mixture of molecules with or without the 2-O-methylfucosyl residue (161). Because of the similarity between *nodZ/nolO* in *B. japonicum* and *nodH/nodPQ* in *R. meliloti*, Luka et al. (161) suggest that NodZ mediates the 2-O-methylfucosylation of the *B. japonicum* Nod metabolites utilizing a NodO-dependent substrate.

R. tropici strains produce a mixture of sulfated and nonsulfated Nod metabolites. Both stimulate external cortical cell division on *Phaseolus vulgaris*, but the sulfated compound is far more active. The Nod metabolites from *R. elii*, another symbiont of *Phaseolus vulgaris*, are quite different from those of *R. tropici*: they are nonsulfated and carry a modified fucose at the reducing end (165). Because of the major differences between the Nod factors of *R. tropici* and *R. elii*, Martinez et al. (165) suggest that *Phaseolus vulgaris* possesses different means to interact with Nod factors.

A. caulinodans also secretes Nod metabolites which are mono-N-acylated chitin oligomers, carrying several unusual substitutions (172). For example, an N-methyl group at the nonreducing end of all the molecules and the reducing glucosamine is substituted by a unique sugar, D-arabinose. Some of the *Azorhizobium* *nod* genes can be implicated in these modifications. The *nodS* gene most probably encodes an S-adenosylmethionine-dependent methyltransferase for Nod factor methylation (80), and a *nolK* gene may be involved in the synthesis of an arabinosyl precursor for factor glycosylation at the reducing end. These Nod metabolites have been found to elicit morphological changes on *Sesbania rostrata* roots: root hair deformation and induction of meristematic loci at lateral root bases, the sites where the root nodules are formed upon bacterial infection (112).

R. leguminosarum bv. *viciae* TOM can efficiently nodulate varieties of peas such as cv. Afghanistan (157). Strain TOM produces Nod factors that are similar to those of other *R. leguminosarum* bv. *viciae* strains. However, one of the nodulation factors made by strain TOM differs in that it carries an O-acetyl group on the C-6 of the reducing N-acetylglucosamine residue. This acetylation is NodX dependent. Although the *nodL* gene product is also an O-acetyltransferase, there is very little similarity between the derived amino acid sequences of the two acetyltransferases (72).

It thus appears that all rhizobia produce Nod metabolites which belong to the same family of molecules, lipooligosaccharides. The common core is probably synthesized by enzymes encoded by the common *nod* genes present in *Rhizobium*, *Bradyrhizobium*, and *Azorhizobium* species. The function of the host-specific *nod* genes is to code for enzymes which decorate the core molecules and make them plant specific.

ROLE OF NodO IN SYMBIOSIS

R. leguminosarum bv. *viciae* *nodE* mutants nodulate *Vicia* plants inefficiently (60). In the absence of the *nodE* gene, the host-specific nodulation factor with an acyl C_{18:4} is not made and only the signal with an acyl C_{18:1} group is formed. This latter signal cannot induce nodule meristem formation when added to *Vicia sativa* plants (245). Deletion analysis revealed that the *nodO* region was responsible for the reduced level of nodulation of peas or vetch observed with *nodE* mutants (60).

The NodO protein shows limited homology to secreted hemolysins and to some proteases (68) and shares a similar secretion mechanism to the secretion of hemolysin and protease (PrtB) by *Erwinia chrysanthemi* (216). Sutton et al. (258) recently showed that the *nodO* gene encodes a secreted protein with no detectable cellulase, pectinase, or protease activity but that it could form Ca^{2+} -regulated ion channels in an artificial membrane, similar to hemolysin-like toxins of certain mammalian pathogens. They concluded, since the pure lipooligosaccharides can induce nodule meristems in the absence of bacteria (245, 263), that NodO must have a complementary role to the nodulation factors. Such a role might be to contribute to the growth of the infection thread by causing a specific ion flux across the plasma membrane. With the recent finding of *nodO* in other rhizobia, such as *Rhizobium* sp. strain BR816 (271) and possibly also *B. japonicum*, this hypothesis can now be explored on a broader basis.

CONCLUSION

Establishment of the *Rhizobium*-plant symbiosis is the result of a multitude of communications between the symbionts and biochemical reactions within the symbionts. The study of mutants has shown that some of the steps can be uncoupled. A great majority of the knowledge is available on the mechanism of nodule formation. Less is known about the biochemistry of nodule function. Nevertheless, it should be realized that the knowledge about *Rhizobium*-plant symbiosis is based on a few selected systems, which might not cover the whole scale of *Rhizobium*-plant symbiosis occurring in nature. From these examples, a general mechanism can be formulated. With the characterization of new *Rhizobium* isolates, it is becoming clear that *Rhizobium* taxonomy requires constant updating. The question can be asked whether these new *Rhizobium* isolates, particularly from leguminous trees, will reveal other molecular mechanisms of nodule formation and/or infection.

In this review, little has been mentioned about plant genes that are essential for the *Rhizobium*-plant symbiosis, despite vast amounts of available information (for reviews, see references 50, 78, 108, and 276). Until now, these studies were based on the isolation and characterization of proteins and their transcripts that are specifically induced in nodule tissue. The proteins are called nodulins. Plant molecular biology now provides new tools to study nodule-specific genes, and it can be expected that application of these techniques (mutagenesis, transposon tagging, transformation, etc.) will shortly provide more knowledge in this area. Already, genetic linkage maps of legumes are being constructed and DNA sequencing techniques are being used to create physical maps of legume genomes (134).

ACKNOWLEDGMENTS

We thank A. Kondorosi, B. J. J. Lugtenberg, R. De Mot, and A. Van Gool for critically reading the manuscript, and we thank A. Vermassen and J. Hansen for their assistance in preparing the manuscript.

P. van Rijn was supported by a grant from Instituut voor Wetenschappelijk Onderzoek in de Nijverheid en de Landbouw.

REFERENCES

- Aguilar, J. M., A. M. Ashby, J. M. Richards, G. J. Loake, M. D. Watson, and C. H. Shaw. 1988. Chemotaxis of *Rhizobium leguminosarum* biovar *phaseoli* towards flavonoid inducers of the symbiotic nodulation genes. *J. Gen. Microbiol.* **134**:2741–2746.
- Allen, O. N., and E. K. Allen. 1981. The Leguminosae. A source book of characteristics, uses and nodulation. University of Wisconsin Press, Madison.
- Appelbaum, E. R., D. V. Thompson, K. Idler, and N. Chartrain. 1988. *Bradyrhizobium japonicum* USDA 191 has two *nodD* genes that differ in

- primary structure and function. *J. Bacteriol.* **170**:12–20.
- Armitage, J. P., A. Gallagher, and A. W. B. Johnston. 1988. Comparison of the chemotactic behaviour of *Rhizobium leguminosarum* with and without the nodulation plasmid. *Mol. Microbiol.* **2**:743–748.
- Arnold, W., A. Becker, M. Keller, A. Roxlau, and A. Pühler. 1994. The role of *Rhizobium meliloti* surface polysaccharides in the infection of *Medicago sativa* nodules. *Endocytobiosis Cell Res.* **10**:17–28.
- Atkinson, E. M., and S. R. Long. 1992. Homology of *Rhizobium meliloti* NodC to polysaccharide polymerizing enzymes. *Mol. Plant-Microbe Interact.* **5**:439–442.
- Atkinson, E. M., M. M. Palcic, O. Hindsgaul, and S. R. Long. 1994. Biosynthesis of *Rhizobium meliloti* lipopolysaccharide Nod factors: NodA is required for an N-acyltransferase. *Proc. Natl. Acad. Sci. USA* **91**:8418–8422.
- Baev, N., G. Endre, G. Petrovics, Z. Bánfalvi, and A. Kondorosi. 1991. Six nodulation genes of *nod* box locus 4 in *Rhizobium meliloti* are involved in nodulation signal production: *nodM* codes for D-glucosamine synthetase. *Mol. Gen. Genet.* **228**:113–124.
- Baev, N., and A. Kondorosi. 1992. Nucleotide sequence of the *R. meliloti* *nodL* gene located in locus n5 of the *nod* regulon. *Plant Mol. Biol.* **18**:843–846.
- Baev, N., M. Schultze, I. Barlier, D. C. Ha, H. Virelizier, E. Kondorosi, and A. Kondorosi. 1992. *Rhizobium nodD* and *nodN* genes are common nod genes: *nodM* encodes functions for efficiency of Nod signal production and bacteroid maturation. *J. Bacteriol.* **174**:7555–7565.
- Bakhuizen, R. 1988. The plant cytoskeleton in the *Rhizobium*-legume symbiosis. Ph.D. thesis. University of Leiden, Leiden, The Netherlands.
- Barbour, W. M., D. R. Hatterman, and G. Stacey. 1991. Chemotaxis of *Bradyrhizobium japonicum* to soybean exudates. *Appl. Environ. Microbiol.* **57**:2635–2639.
- Barnett, M. J., and S. R. Long. 1990. DNA sequence and translational product of a new nodulation-regulatory locus: SyrM has sequence similarity to NodD proteins. *J. Bacteriol.* **172**:3695–3700.
- Barry, M. A., and J. A. Downie. 1993. Identification of the NodC protein in the inner but not the outer membrane of *Rhizobium leguminosarum*. *Mol. Plant-Microbe Interact.* **8**:669–672.
- Bec-Ferté, M. P., H. B. Krishnan, D. Promé, A. Savagnac, S. G. Pueppke, and J. C. Promé. 1994. Structures of nodulation factors from the nitrogen-fixing soybean symbiont *Rhizobium fredii* USDA257. *Biochemistry* **33**:11782–11788.
- Bhuvaneswari, T. V., A. A. Bhagwat, and W. D. Bauer. 1981. Transient susceptibility of root cells in four common legumes: its nodulation of rhizobia. *Plant Physiol.* **68**:1144–1149.
- Bhuvaneswari, T. V., and B. Solheim. 1985. Root hair deformation in the white clover/*Rhizobium trifolii* symbiosis. *Physiol. Plant.* **63**:25–31.
- Bibb, M. J., S. Biro, H. Motamedi, J. F. Collins, and C. R. Hutchinson. 1989. Analysis of the nucleotide sequence of the *Streptomyces glaucescens* *tcml* genes provide key information about the enzymology of polyketide antibiotic biosynthesis. *EMBO J.* **9**:2727–2736.
- Bloemberg, G. V., J. E. Thomasoates, B. J. J. Lugtenberg, and H. P. Spaik. 1994. Nodulation protein NodL of *Rhizobium leguminosarum* O-acetylates lipo-oligosaccharides, chitin fragments and N-acetylglucosamine in vitro. *Mol. Microbiol.* **11**:793–804.
- Bohloul, B. B., and E. L. Schmidt. 1974. Lectins: a possible basis for specificity in the *Rhizobium*-legume root nodule symbiosis. *Science* **185**:269–271.
- Boundy-Mills, K. L., R. M. Kossak, R. E. Tully, S. G. Pueppke, S. Lohrke, M. J. Sadowsky. 1994. Induction of the *Rhizobium fredii* *nod*-box independent nodulation gene *nodJ* requires a functional *nodD1* gene. *Mol. Plant-Microbe Interact.* **7**:305–308.
- Breedveld, M. W., and K. J. Miller. 1994. Cyclic β -glucans of members of the family of *Rhizobiaceae*. *Microbiol. Rev.* **58**:145–161.
- Brewin, N. J. 1991. Development of the legume root nodule. *Annu. Rev. Cell Biol.* **7**:191–226.
- Broughton, W. J., A. Krause, A. Lewin, X. Perret, and N. Price. 1991. Signal exchange mediates host-specific nodulation of tropical legumes by the broad-host-range *Rhizobium* species NGR234, p. 162–167. In H. Hennecke, and D. P. S. Verma (ed.), *Advances in molecular genetics of plant-microbe interactions*, vol. 1. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Bulawa, C. E., and W. Wasco. 1991. Chitin and nodulation. *Nature (London)* **353**:710.
- Burn, J., L. Rossen, and A. W. B. Johnston. 1987. Four classes of mutations in the *nodD* gene of *Rhizobium leguminosarum* biovar *viciae* that affects its ability to autoregulate and/or activate other *nod* genes in the presence of flavonoid inducers. *Genes Dev.* **1**:456–464.
- Burn, J. E., W. D. Hamilton, J. C. Wootton, and A. W. B. Johnston. 1989. Single and multiple mutations affecting properties of the regulatory gene *nodD* of *Rhizobium*. *Mol. Microbiol.* **3**:1567–1577.
- Caetano-Anollés, G., D. K. Crist-Estes, and W. D. Bauer. 1988. Chemotaxis of *Rhizobium meliloti* to the plant flavone luteolin requires functional nodulation genes. *J. Bacteriol.* **170**:3164–3169.

28. Caetano-Anollés, G., E. Wrobel-Boerner, and W. D. Bauer. 1992. Growth and movement of spot inoculated *Rhizobium meliloti* on the root surface of alfalfa. *Plant Physiol.* **98**:1181–1189.
29. Callaham, D. A., and J. G. Torrey. 1981. The structural basis for the infection of root hairs in *Trifolium repens* by *Rhizobium*. *Can. J. Bot.* **59**: 1647–1664.
30. Calvert, H. E., M. K. Pence, M. Pierce, N. S. A. Malik, and W. Bauer. 1984. Anatomical analysis of the development and distribution of *Rhizobium* infection in soybean roots. *Can. J. Bot.* **62**:2375–2384.
31. Carlson, R. W., S. J. Juan, U. R. Bhat, J. Glushka, H. P. Spink, A. H. M. Wijffes, A. N. N. van Brussel, T. J. W. Stokkermans, N. K. Peters, and G. Stacey. 1993. The structures and biological activities of the lipo-oligosaccharide nodulation signals produced by type-1 and type-2 strains of *Bradyrhizobium japonicum*. *J. Biol. Chem.* **268**:18372–18381.
32. Chandler, M. R. 1978. Some observations on the infection of *Arachis hypogaea* L. by *Rhizobium*. *J. Exp. Bot.* **29**:749–755.
33. Chandler, M. R., R. A. Date, and R. J. Roughley. 1982. Infection and root nodule development in *Stylosanthes* species by *Rhizobium*. *J. Exp. Bot.* **33**:47–57.
34. Chua, Y.-K., C. E. Pankhurst, P. E. Macdonald, D. Hopcroft, B. D. W. Jarvis, and D. B. Scott. 1985. Isolation and characterisation of Tn5-induced exudates contain elevated levels of daidzein and coumestrol in response to *Rhizobium* inoculation. *Mol. Plant-Microbe Interact.* **6**:665–668.
35. Cooper, E. J., and J. R. Rao. 1992. Localized changes in flavonoid biosynthesis in roots of *Lotus pedunculatus* after infection by *Rhizobium loti*. *Plant Physiol.* **100**:444–450.
36. Cooper, J. B., and S. R. Long. 1994. Morphogenetic rescue of *Rhizobium meliloti* nodulation mutants by *trans*-zeatin secretion. *Plant Cell* **6**:215–225.
37. Cren, M., A. Kondorosi, and E. Kondorosi. 1993. An insertional point mutation inactivates NodR repressor in *Rhizobium meliloti* 1021. *J. Bacteriol.* **176**:518–519.
38. Dakora, F. D., C. M. Joseph, and D. A. Phillips. 1993. Alfalfa (*Medicago sativa* L.) root exudates contain isoflavonoids in the presence of *Rhizobium meliloti*. *Plant Physiol.* **101**:819–824.
39. Dakora, F. D., C. M. Joseph, and D. A. Phillips. 1993. Common bean root exudates contain elevated levels of daidzein and coumestrol in response to *Rhizobium* inoculation. *Mol. Plant-Microbe Interact.* **6**:665–668.
40. Dart, P. 1977. Infection and development of leguminous nodules, p. 367–472. In R. W. F. Hardy and W. S. Silver (ed.), *A treatise on dinitrogen fixation*. John Wiley & Sons, Inc., New York.
41. Davis, E. O., I. J. Evans, and A. W. B. Johnston. 1988. Identification of *nodX*, a gene that allows *Rhizobium leguminosarum* biovar *viciae* strain TOM to nodulate Afghanistan peas. *Mol. Gen. Genet.* **212**:531–535.
42. Davis, E. O., and A. W. B. Johnston. 1990. Analysis of three *nodD* genes of *Rhizobium leguminosarum* bv. *phaseoli*: *nodD₁* is preceded by *nolE*, a gene whose product is secreted from the cytoplasm. *Mol. Microbiol.* **4**:921–932.
43. Davis, E. O., and A. W. B. Johnston. 1990. Regulatory functions of the 3 *nodD* genes of *Rhizobium leguminosarum* bv. *phaseoli*. *Mol. Microbiol.* **4**:933–941.
44. de Faria, S. M., H. T. Hay, and J. I. Sprent. 1988. Entry of rhizobia into roots of *Mimosa scabrella* Benth occurs between epidermal cells. *J. Gen. Microbiol.* **134**:2291–2296.
45. de Faria, S. M., G. P. Lewis, J. I. Sprent, and J. M. Sutherland. 1989. Occurrence of nodulation in the leguminosae. *New Phytol.* **111**:607–619.
46. de Faria, S. M., S. G. McIlmoy, and J. I. Sprent. 1987. The occurrence of infected cells with persistent infection threads in legume root nodules. *Can. J. Bot.* **65**:553–558.
47. de Faria, S. M., J. M. Sutherland, and J. I. Sprent. 1986. A new type of infected cell in root nodules of *Andira* spp. (leguminosae). *Plant Sci.* **45**: 143–147.
48. Debellé, F., C. Rosenberg, and J. Dénarié. 1992. The *Rhizobium*, *Bradyrhizobium*, and *Azorhizobium* NodC proteins are homologous to yeast chitin synthases. *Mol. Plant-Microbe Interact.* **5**:443–446.
49. Debellé, F., and S. B. Sharma. 1986. Nucleotide sequence of *Rhizobium meliloti* RCR2011 genes involved in host specificity of nodulation. *Nucleic Acids Res.* **14**:7451–7472.
50. De Bruijn, F. J., R. J. Chen, S. Y. Fujimoto, A. Pinaev, D. Silver, and K. Szczygłowski. 1994. Regulation of nodulin gene expression. *Plant Soil* **161**:59–68.
51. Demont, N., M. Ardourel, F. Maillet, D. Promé, M. Ferro, J.-C. Promé, and J. Dénarié. 1994. The *Rhizobium meliloti* regulatory *nodD3* and *syrM* genes control the synthesis of a particular class of nodulation factors N-acetylated by (omega-1)-hydroxylated fatty acids. *EMBO J.* **13**:2139–2149.
52. Demont, N., F. Debellé, H. Aurelle, J. Dénarié, and J. C. Promé. 1993. Role of the *Rhizobium meliloti* *nodF* and *nodE* genes in the biosynthesis of lipo-oligosaccharide nodulation factors. *J. Biol. Chem.* **268**:20143–20142.
- 52a. Dénarié, J., and J. Cullimore. 1993. Lipo-oligosaccharide nodulation factors: a new class of signalling molecules mediating recognition and morphogenesis. *Cell* **74**:951–954.
53. Dénarié, J., F. Debellé, and C. Rosenberg. 1992. Signaling and host range variation in nodulation. *Annu. Rev. Microbiol.* **46**:497–531.
- 53a. Diaz, C. L., L. S. Melchers, P. J. J. Hooykaas, B. J. J. Lugtenberg, and J. W. Kijne. 1989. Root lectin as a determinant of host-plant specificity in the *Rhizobium* legume symbiosis. *Nature* (London) **338**:579–581.
54. Djordjevic, M. A., J. W. Redmond, M. Batley, and B. G. Rolfe. 1987. Clovers secrete specific phenolic compounds which either stimulate or repress *nod* gene expression in *Rhizobium trifolii*. *EMBO J.* **6**:1173–1179.
55. Dockendorff, T. C., A. J. Sharma, and G. Stacey. 1994. Identification and characterization of the *nolYZ* genes of *Bradyrhizobium japonicum*. *Mol. Plant-Microbe Interact.* **7**:173–180.
56. Dowling, D. N., and W. J. Broughton. 1986. Competition for nodulation of legumes. *Annu. Rev. Microbiol.* **40**:131–157.
57. Downie, J. A. 1989. The *nodL* gene from *Rhizobium leguminosarum* is homologous to the acetyltransferases encoded by *lacA* and *cysE*. *Mol. Microbiol.* **3**:1649–1651.
58. Downie, J. A. 1991. A *nod* of recognition. *Curr. Biol.* **1**:382–384.
59. Downie, J. A., C. D. Knight, A. W. B. Johnston, and L. Rossen. 1985. Identification of genes and gene products involved in the nodulation of peas by *Rhizobium leguminosarum*. *Mol. Gen. Genet.* **198**:255–262.
60. Downie, J. A., and B. P. Surin. 1990. Either of two *nod* gene loci can complement the nodulation defect of a nod deletion mutant of *Rhizobium leguminosarum* bv. *viciae*. *Mol. Gen. Genet.* **222**:81–86.
61. Dreyfus, B. L., J. L. Garcia, and M. Gillis. 1988. Characterization of *Azorhizobium caulinodans* gen. nov., sp. nov., a stem-nodulating nitrogen-fixing bacterium isolated from *Sesbania rostrata*. *Int. J. Syst. Bacteriol.* **38**:89–98.
62. Dreyfus, B. L., and Y. R. Dommergues. 1981. Nitrogen-fixing nodules induced by *Rhizobium* on the stem of the tropical legume *Sesbania rostrata*. *FEMS Microbiol. Lett.* **10**:313–317.
63. Dudley, M. E., T. W. Jacobs, and S. R. Long. 1987. Microscopy studies of cell division induced in alfalfa roots by *Rhizobium meliloti*. *Planta* **171**:289–301.
64. Dudman, W. F. 1984. The polysaccharides and oligosaccharides of *Rhizobium* and their role in the infection process, p. 397–404. In C. Veeger and W. E. Newton (ed.), *Advances in nitrogen fixation research*. Martinus Nijhoff, The Hague, The Netherlands.
65. Duhoux, E. 1984. Ontogénèse des nodules caulinaires de *Sesbania rostrata* (légumineuse). *Can. J. Bot.* **62**:982–994.
66. Dusha, I., A. Bakos, A. Kondorosi, F. J. de Bruijn, and J. Schell. 1989. The *Rhizobium meliloti* early nodulation genes (*nodABC*) are nitrogen-regulated: isolation of a mutant strain with efficient nodulation capacity of alfalfa in the presence of ammonium. *Mol. Gen. Genet.* **219**:89–96.
67. Eardly, B. D., D. B. Hannaway, and P. J. Bottemley. 1985. Characterization of rhizobia from ineffective alfalfa nodules: ability to nodulate bean plants *Phaseolus vulgaris* (L.). *Appl. Environ. Microbiol.* **50**:1422–1427.
68. Economou, A., W. D. O. Hamilton, A. W. B. Johnston, and J. A. Downie. 1990. The *Rhizobium* nodulation gene *nodO* encodes a Ca²⁺-binding protein that is exported without N-terminal cleavage and is homologous to haemolysin and related proteins. *EMBO J.* **9**:349–354.
69. Eskew, D. L., Q. Y. Jiang, G. Caetano-Anolles, and P. M. Gresshoff. 1993. Kinetics of nodule development in *Glycine* soja. *Plant Physiol.* **103**:1139–1145.
70. Evans, I. J., and J. A. Downie. 1986. The NodI product of *Rhizobium leguminosarum* is closely related to ATP-binding bacterial transport proteins: nucleotide sequence of the *nodI* and *nodJ* genes. *Gene* **43**:95–101.
71. Faucher, C., H. Camut, J. Dénarié, and G. Truchet. 1989. The *nodH* and *nodQ* host range genes of *Rhizobium meliloti* behave as avirulence genes in *R. leguminosarum* bv. *viciae* and determine changes in the production of plant-specific extracellular signals. *Mol. Plant-Microbe Interact.* **2**:291–300.
72. Firmin, J. L., K. E. Wilson, R. W. Carlson, A. E. Davies, and J. A. Downie. 1993. Resistance to nodulation of cv Afghanistan peas is overcome by *nodX*, which mediates an O-acetylation of the *Rhizobium leguminosarum* lipo-oligosaccharide nodulation factor. *Mol. Microbiol.* **10**:351–360.
73. Firmin, J. L., K. E. Wilson, L. Rossen, and A. W. B. Johnston. 1986. Flavonoid activation of nodulation genes in *Rhizobium* reversed by other compounds present in plants. *Nature* (London) **324**:90–92.
74. Fisher, R. F., T. T. Egelhoff, J. T. Mulligan, and S. R. Long. 1988. Specific binding of proteins from *Rhizobium meliloti* cell-free extracts containing NodD to DNA sequences upstream of inducible nodulation genes. *Genes Dev.* **2**:282–293.
75. Fisher, R. F., and S. R. Long. 1992. *Rhizobium*-plant signal exchange. *Nature* (London) **357**:655–660.
76. Fisher, R. F., and S. R. Long. 1993. Interactions of NodD at the *nod* box: NodD binds to two distinct sites on the same face of the helix and induces a bend in the DNA. *J. Mol. Biol.* **233**:336–348.
77. Fisher, R. F., J. A. Swanson, J. T. Mulligan, and S. R. Long. 1987. Extended region of nodulation genes in *Rhizobium meliloti* 1021. II. Nucleotide sequence, transcription start sites and protein products. *Genetics* **117**:191–201.
78. Franssen, H. J., I. Vijn, W. C. Yang, and T. Bisseling. 1992. Developmental aspects of the *Rhizobium*-legume symbiosis. *Plant Mol. Biol.* **19**:89–107.
79. Gaworzewska, E. T., and M. J. Carlile. 1982. Positive chemotaxis of *Rhizobium leguminosarum* and other bacteria towards root exudates from legumes and other plants. *J. Gen. Microbiol.* **128**:789–798.
80. Geelen, D., P. Mergaert, R. A. Geremia, S. Goormachtig, M. Van Montagu,

- and M. Holsters. 1993. Identification of *nodSUII* genes on Nod locus 1 of *Azorhizobium caulinodans*: evidence that *nodS* encodes a methyltransferase involved in Nod factor modification. *Mol. Microbiol.* **9**:145–154.
81. Geiger, O., H. P. Spaink, and E. P. Kennedy. 1991. Isolation of *Rhizobium leguminosarum* NodF nodulation protein: NodF carries a 4' phosphopantetheine prosthetic group. *J. Bacteriol.* **173**:2872–2878.
 82. Geiger, O., J. E. Thomasates, J. Glushka, H. P. Spaink, and B. J. J. Lugtenberg. 1994. Phospholipids of *Rhizobium* contain *nodE*-determined highly unsaturated fatty acid moieties. *J. Biol. Chem.* **269**:11090–11097.
 83. Geremia, R. A., P. Mergaert, D. Geelen, M. Van Montagu, and M. Holsters. 1994. The NodC protein of *Azorhizobium caulinodans* is an N-acetylglucosaminyltransferase. *Proc. Natl. Acad. Sci. USA* **91**:2669–2673.
 84. Gibson, A. 1968. Nodulation failure in *Trifolium subterraneum* L. cv. Woonenellup (Syn. Marrar.). *Aust. J. Agric. Sci.* **19**:907–918.
 85. Goethals, K., M. Gao, K. Tomeke, M. Van Montagu, and M. Holsters. 1989. Common *nodABC* genes in nod locus 1 of *Rhizobium caulinodans*: nucleotide sequence and plant-inducible expression. *Mol. Gen. Genet.* **219**:289–298.
 86. Goethals, K., P. Mergaert, M. Gao, D. Geelen, M. Van Montagu, and M. Holsters. 1992. Identification of new inducible nodulation genes in *Azorhizobium caulinodans*. *Mol. Plant-Microbe Interact.* **5**:405–411.
 87. Goethals, K., G. Van Den Eede, M. Van Montagu, and M. Holsters. 1990. Identification and characterization of a functional *nodD* gene in *Azorhizobium caulinodans* ORS571. *J. Bacteriol.* **172**:2658–2666.
 88. Goethals, K., M. Van Montagu, and M. Holsters. 1992. Conserved motifs in a divergent *nod* box of *Azorhizobium caulinodans* ORS571 reveal a common structure in promoters regulated by LysR-type proteins. *Proc. Natl. Acad. Sci. USA* **89**:1646–1650.
 89. Göttfert, M., P. Grob, and H. Hennecke. 1990. Proposed regulatory pathway encoded by the *nodV* and *nodW* genes, determinants of host specificity in *Bradyrhizobium japonicum*. *Proc. Natl. Acad. Sci. USA* **87**:2680–2684.
 90. Göttfert, M., S. Hitz, and H. Hennecke. 1990. Identification of *nodS* and *nodU*, two inducible genes inserted between the *Bradyrhizobium japonicum* *nodYAB* and *nodIJ* genes. *Mol. Plant-Microbe Interact.* **3**:308–316.
 91. Göttfert, M., D. Holzäuser, D. Bani, and H. Hennecke. 1992. Structural and functional analysis of two different *nodD* genes in *Bradyrhizobium japonicum* USDA110. *Mol. Plant-Microbe Interact.* **5**:257–265.
 92. Göttfert, M., B. Horvath, E. Kondorosi, P. Putnoky, F. Rodrigues-Quinones, and A. Kondorosi. 1986. At least two *nodD* genes are necessary for efficient nodulation of alfalfa by *Rhizobium meliloti*. *J. Mol. Biol.* **191**:411–420.
 93. Göttfert, M., J. W. Lamb, R. Grasser, J. Semenza, and H. Hennecke. 1989. Mutational analysis of the *Bradyrhizobium japonicum* common *nod* genes and further *nod* box-linked genomic DNA regions. *Mol. Gen. Genet.* **215**:407–415.
 94. Graham, T. L. 1991. Flavonoid and isoflavonoid distribution in developing soybean seedling tissues and in seed and root exudates. *Plant Physiol.* **95**:594–603.
 95. Gray, J. X., and B. G. Rolfe. 1990. Exopolysaccharide production in *Rhizobium* and its role in invasion. *Mol. Microbiol.* **4**:1425–1431.
 96. Grob, P., H. Hennecke, and M. Gottfert. 1993. A novel response-regulator is able to suppress the nodulation defect of a *Bradyrhizobium japonicum* *nodW* mutant. *Mol. Gen. Genet.* **241**:531–541.
 97. Grob, P., H. Hennecke, and M. Gottfert. 1994. Cross-talk between the two-component regulatory systems NodVW and NwsAB of *Bradyrhizobium japonicum*. *FEMS Microbiol. Lett.* **120**:349–354.
 98. Gulash, M., P. Ames, R. C. LaRosiere, and K. Bergman. 1984. Rhizobia are attracted to localized sites on legume roots. *Appl. Environ. Microbiol.* **48**:149–152.
 99. Györgypal, Z., N. Iyer, and A. Kondorosi. 1988. Three regulatory *nodD* alleles of diverged flavonoid-specificity are involved in host-dependent nodulation mutants of *Rhizobium meliloti*. *Mol. Gen. Genet.* **212**:85–92.
 100. Györgypal, Z., and A. Kondorosi. 1991. Homology of the ligand-binding regions of *Rhizobium* symbiotic regulatory protein NodD and vertebrate nuclear receptors. *Mol. Gen. Genet.* **226**:337–340.
 101. Györgypal, Z., E. Kondorosi, and A. Kondorosi. 1991. Diverse signal sensitivity of NodD protein homologs from narrow and broad host range rhizobia. *Mol. Plant-Microbe Interact.* **4**:356–364.
 102. Hamblin, J., and S. P. Kent. 1973. Possible role of phytohemagglutinin in *Phaseolus vulgaris* L. *Nature (London)* **254**:28–30.
 103. Hartwig, U. A., C. A. Maxwell, C. M. Joseph, and D. A. Phillips. 1989. Interactions among flavonoid *nod* gene inducers released from alfalfa seeds and roots. *Plant Physiol.* **91**:1138–1142.
 104. Hartwig, U. A., C. A. Maxwell, C. M. Joseph, and D. A. Phillips. 1990. Chrysoeriol and luteolin released from alfalfa seeds induce *nod* genes in *Rhizobium meliloti*. *Plant Physiol.* **92**:116–122.
 105. Hartwig, U. A., C. A. Maxwell, C. M. Joseph, and D. A. Phillips. 1990. Effects of alfalfa *nod* gene-inducing flavonoids on *nodABC* transcription in *Rhizobium meliloti* strains containing different *nodD* genes. *J. Bacteriol.* **172**:2769–2773.
 - 105a. Heidstra, R., R. Geurts, H. Franssen, H. P. Spaink, A. van Kammen, and T. Bisseling. 1994. Root hair deformation activity of nodulation factors and their fate in *Vicia sativa*. *Plant Physiol.* **105**:787–797.
 106. Henikoff, S., G. W. Haughn, J. M. Calvo, and J. C. Wallace. 1988. A large family of bacterial activator proteins. *Proc. Natl. Acad. Sci. USA* **85**:6602–6606.
 107. Heron, D. S. T., and S. G. Pueppke. 1984. Mode of infection, nodulation specificity, and indigenous plasmids of 11 fast-growing *Rhizobium japonicum* strains. *J. Bacteriol.* **160**:1061–1066.
 108. Hirsch, A. M. 1992. Developmental biology of legume nodulation. *New Phytol.* **122**:211–237.
 109. Hirsch, A. M., T. V. Bhuvaneswari, J. G. Torrey, and T. Bisseling. 1989. Early nodulin genes are induced in alfalfa root outgrowths elicited by auxin transport inhibitors. *Proc. Natl. Acad. Sci. USA* **86**:1244–1248.
 110. Ho, S. C., M. Schindler, and J. L. Wang. 1990. Carbohydrate binding activities of *Bradyrhizobium japonicum*. II. Isolation and characterization of a galactose-specific lectin. *J. Cell Biol.* **111**:1639–1643.
 111. Ho, S. C., J. L. Wang, and M. Schindler. 1990. Carbohydrate binding activities of *Bradyrhizobium japonicum*. I. Saccharide-specific inhibition of homotypic and heterotypic adhesion. *J. Cell Biol.* **111**:1631–1638.
 112. Holsters, M., D. Geelen, K. Goethals, M. Van Montagu, R. Geremia, J. C. Promé, and P. Mergaert. 1993. Nod factor production by *Azorhizobium caulinodans* strain ORS571, p. 191–196. In R. Palacios, J. Mora, and W. E. Newton (ed.), *New horizons in nitrogen fixation*. Kluwer Academic Publishers, Dordrecht, The Netherlands.
 113. Hong, G. F., J. E. Burn, and A. W. B. Johnston. 1987. Evidence that DNA involved in the expression of nodulation (*nod*) genes in *Rhizobium* binds to the product of the regulatory gene *nodM*. *Nucleic Acids Res.* **15**:9677–9690.
 114. Honma, M. A., M. Asomaning, and F. M. Ausubel. 1990. *Rhizobium meliloti* *nodD* genes mediate host-specific activation of *nodABC*. *J. Bacteriol.* **172**:901–911.
 115. Honma, M. A., and F. M. Ausubel. 1987. *Rhizobium meliloti* has three functional copies of the *nodD* symbiotic regulatory gene. *Proc. Natl. Acad. Sci. USA* **84**:8558–8562.
 116. Horvath, B., C. W. Bachem, J. Schell, and A. Kondorosi. 1987. Host-specific regulation of nodulation genes in *Rhizobium* is mediated by a plant signal interacting with the *nodD* gene product. *EMBO J.* **6**:841–848.
 117. Horvath, B., R. Heidstra, M. Lados, M. Moerman, H. P. Spaink, J. C. Promé, A. van Kammen, and T. Bisseling. 1993. Lipo-oligosaccharides of *Rhizobium* induce infection-related early nodulin gene expression in pea root hairs. *Plant J.* **4**:727–733.
 118. Horvath, B., E. Kondorosi, M. John, J. Schmidt, I. Török, Z. Györgypal, I. Barabas, U. Wieneke, J. Schell, and A. Kondorosi. 1986. Organization, structure and symbiotic function of *Rhizobium meliloti* nodulation genes determining host specificity for alfalfa. *Cell* **46**:335–343.
 119. Hubac, C., J. Ferran, D. Guerrier, A. Tremolieres, and A. Kondorosi. 1993. Luteolin absorption in *Rhizobium meliloti* wild-type and mutant strains. *J. Gen. Microbiol.* **139**:1571–1578.
 120. Hungria, M., C. M. Joseph, and D. A. Phillips. 1991. Anthocyanidins and flavonols: major *nod* gene inducers from seeds of a black-seeded common bean (*Phaseolus vulgaris* L.). *Plant Physiol.* **97**:751–758.
 121. Hungria, M., C. M. Joseph, and D. A. Phillips. 1991. *Rhizobium nod* gene inducers exuded naturally from roots of common bean (*Phaseolus vulgaris* L.). *Plant Physiol.* **97**:759–764.
 122. Innes, R. W., P. L. Kuempel, J. Plazinski, H. Canter-Cremers, B. G. Rolfe, and M. A. Djordjevic. 1985. Plant factors induce expression of nodulation and host-range genes in *Rhizobium trifolii*. *Mol. Gen. Genet.* **201**:426–432.
 123. Jacobs, T. W., T. T. Egelhoff, and S. R. Long. 1985. Physical and genetic map of a *Rhizobium meliloti* gene region and nucleotide sequence of *nodC*. *J. Bacteriol.* **162**:469–476.
 124. Jarvis, B. D. W., M. Gillis, and J. De Ley. 1986. Intra- and intergeneric similarities between the ribosomal ribonucleic acid cistrons of *Rhizobium* and *Bradyrhizobium* species and some related bacteria. *Int. J. Syst. Bacteriol.* **36**:129–138.
 125. John, M., H. Röhrig, J. Schmidt, U. Wieneke, and J. Schell. 1993. Rhizobium nodB protein involved in nodulation signal synthesis is a chito oligosaccharide deacetylase. *Proc. Natl. Acad. Sci. USA* **90**:625–629.
 126. Johnson, D., E. L. Roth, and G. Stacey. 1989. Immunogold localization of the NodC and NodA proteins of *Rhizobium meliloti*. *J. Bacteriol.* **171**:4583–4588.
 127. Jordan, D. C. 1984. Family III. *Rhizobiaceae* Conn 1938, 321^{AL}, p. 234–254. In N. R. Krieg and J. G. Holt (ed.), *Bergey's manual of systematic bacteriology*, vol. 1. The Williams & Wilkins Co., Baltimore.
 128. Journet, J. P., N. Pichon, A. Dedieu, F. de Billy, G. Truchet, and D. G. Barker. 1994. *Rhizobium meliloti* Nod factors elicit cell-specific transcription of the *Enod12* gene in transgenic alfalfa. *Plant J.* **6**:241–249.
 129. Kape, R., M. Parniske, and D. Werner. 1991. Chemotaxis and *nod* gene activity of *Bradyrhizobium japonicum* in response to hydrocinnamic acids and isoflavonoids. *Appl. Environ. Microbiol.* **57**:316–319.
 130. Keyser, H. H., B. B. Bohlool, T. S. Hu, and D. F. Weber. 1982. Fast-growing rhizobia isolated from roots of soybean. *Science* **215**:1631–1632.
 131. Kijne, J. W. 1975. The fine structure of pea root nodule. 1. Vacuolar changes after endocytotic host cell infection by *Rhizobium leguminosarum*. *Physiol. Plant Pathol.* **5**:75–79.
 132. Kijne, J. W. 1992. The *Rhizobium* infection process, p. 348–397. In G.

- Stacey, R. H. Burris, and H. J. Evans (ed.), Biological nitrogen fixation. Chapman & Hall, New York.
133. Kijne, J. W., G. Smit, C. L. Diaz, and B. J. J. Lugtenberg. 1988. Lectin-enhanced accumulation of manganese-limited *Rhizobium leguminosarum* cells on pea root hair tips. *J. Bacteriol.* **170**:2994–3000.
 134. Kiss, G. B., G. Csanadi, K. Kalman, P. Kalo, and L. Okresz. 1993. Construction of a basic genetic map for alfalfa using RFLP, RAPD, isozyme and morphological markers. *Mol. Gen. Genet.* **238**:129–137.
 135. Kondorosi, A. 1991. Overview on genetics of nodule induction: factors controlling nodule induction by *Rhizobium meliloti*, p. 111–118. In H. Hennecke and D. P. S. Verma (ed.), *Advances in molecular genetics of plant-microbe interactions*, vol. 1. Kluwer Academic Publishers, Dordrecht, The Netherlands.
 136. Kondorosi, A. 1992. Regulation of nodulation genes in rhizobia, p. 325–340. In D. P. S. Verma (ed.), *Molecular signals in plant-microbe communication*. CRC Press, Inc., Boca Raton, Fla.
 137. Kondorosi, E., Z. Banfalvi, and A. Kondorosi. 1984. Physical and genetic analysis of a symbiotic region of *Rhizobium meliloti*: identification of nodulation genes. *Mol. Gen. Genet.* **193**:445–452.
 138. Kondorosi, E., M. Buiré, M. Cren, N. Iyer, B. Hoffmann, and A. Kondorosi. 1991. Involvement of the *sydM* and *nodD3* genes of *Rhizobium meliloti* in *nod* gene activation and in optimal nodulation of the plant host. *Mol. Microbiol.* **5**:3035–3048.
 139. Kondorosi, E., J. Gyuris, J. Schmidt, M. John, E. Duda, B. Hoffman, J. Schell, and A. Kondorosi. 1989. Positive and negative control of *nod* gene expression in *Rhizobium meliloti* is required for optimal nodulation. *EMBO J.* **5**:1331–1340.
 140. Kondorosi, E., M. Pierre, M. Cren, U. Haumann, M. Buiré, B. Hoffman, J. Schell, and A. Kondorosi. 1991. Identification of *NolR*, a negative trans-acting factor controlling the *nod* regulon in *Rhizobium meliloti*. *J. Mol. Biol.* **222**:885–896.
 141. Kossiak, R. M., R. S. Joshi, B. A. Bowen, H. E. Paaren, and E. R. Appelbaum. 1990. Strain-specific inhibition of *nod* gene induction in *Bradyrhizobium japonicum* by flavonoid compounds. *Appl. Environ. Microbiol.* **56**:1333–1341.
 142. Kredich, N. M. 1987. Biosynthesis of cysteine, p. 419–428. In F. C. Neidhardt, J. L. Ingraham, K. B. Low, B. Magasanik, M. Schaechter, and H. E. Umbarger (ed.), *Escherichia coli* and *Salmonella typhimurium*: cellular and molecular biology. American Society for Microbiology, Washington, D.C.
 143. Krishnan, H. B., A. Lewin, R. Fellay, W. J. Broughton, and S. G. Pueppke. 1992. Differential expression of *nodS* accounts for the varied abilities of *Rhizobium fredii* USDA257 and *Rhizobium* sp. NGR234 to nodulate *Leucaena* spp. *Mol. Microbiol.* **6**:3321–3330.
 144. Krishnan, H. B., and S. G. Pueppke. 1991. *nolC*, a *Rhizobium fredii* gene involved in cultivar specific nodulation of soybean, shares homology with a heat-shock gene. *Mol. Microbiol.* **5**:737–745.
 145. Krishnan, H. B., and S. G. Pueppke. 1992. Inactivation of *nolC* conditions developmental abnormalities in nodulation of Peking soybean by *Rhizobium fredii* USDA257. *Mol. Plant-Microbe Interact.* **5**:14–21.
 146. Krishnan, H. B., and S. G. Pueppke. 1994. Cultivar-specificity genes of the nitrogen-fixing soybean symbiont, *Rhizobium fredii* USDA257, also regulate nodulation of *Erythrina* spp. *Am. J. Bot.* **8**:38–45.
 147. Lancelle, S. A., and J. G. Torrey. 1984. Early development of R-induced root nodules of *Parasponia rigida*. I. Infection and early nodule initiation. *Protoplasma* **123**:26–37.
 148. Leigh, J. A., and D. L. Coplin. 1992. Exopolysaccharides in plant-bacterial interactions. *Annu. Rev. Microbiol.* **46**:307–346.
 149. Leigh, J. A., and G. C. Walker. 1994. Exopolysaccharides of *Rhizobium*—synthesis, regulation and symbiotic function. *Trends Genet.* **10**:63–67.
 150. Lerouge, P., P. Roche, C. Faucher, F. Maillet, G. Truchet, J. C. Promé, and J. Dénarié. 1990. Symbiotic host-specificity of *Rhizobium meliloti* is determined by sulphated and acylated glucosamine oligosaccharide. *Nature (London)* **344**:781–784.
 151. Le Strange, K. K., G. L. Bender, M. A. Djordjevic, B. G. Rolfe, and J. W. Redmond. 1990. The *Rhizobium* strain NGR234 *nodD1* gene product responds to activation by simple phenolic compounds vanillin and isovanillin present in wheat seedling extracts. *Mol. Plant-Microbe Interact.* **3**:214–220.
 152. Lewin, A., E. Cervantes, C. H. Wong, and W. J. Broughton. 1990. *nodSU*, two new *nod* genes of the broad host-range *Rhizobium* strain NGR234 encode host-specific nodulation of the tropical tree *Leucaena leucocephala*. *Mol. Plant-Microbe Interact.* **3**:317–326.
 153. Lewin, A., C. Rosenberg, Z. A. H. Meyer, C. H. Wong, L. Nelson, J. F. Manen, J. Stanley, D. N. Downing, J. Dénarié, and W. J. Broughton. 1987. Multiple host-specificity loci of the broad host-range *Rhizobium* sp. NGR234 selected using the widely compatible legume *Vigna unguiculata*. *Plant. Mol. Biol.* **8**:447–459.
 154. Lewis-Henderson, W. R., and M. A. Djordjevic. 1991. A cultivar-specific interaction between *Rhizobium leguminosarum* bv. *trifolii* and subterranean clover is controlled by *nodM*, other bacterial cultivar specificity genes, and a single recessive host gene. *J. Bacteriol.* **173**:2791–2799.
 155. Lewis-Henderson, W. R., and M. A. Djordjevic. 1991. *nodT*, a positively-acting cultivar specificity determinant controlling nodulation of *Trifolium subterraneum* by *Rhizobium leguminosarum* biovar *trifolii*. 1991. *Plant Mol. Biol.* **16**:515–526.
 156. Libbenga, K. R., and P. A. A. Harkes. 1973. Initial proliferation of cortical cells in the formation of root nodules in *Pisum sativum*. *Planta* **114**:17–28.
 157. Lie, T. A. 1978. Symbiotic specialization in pea plants: the requirement of specific *Rhizobium* strains for peas from Afghanistan. *Ann. Appl. Biol.* **88**:462–465.
 158. Liu, R. L., V. M. Tran, and E. L. Schmidt. 1989. Nodulation competitiveness of a nonmotile Tn7-mutant of *Bradyrhizobium japonicum* in nonsterile soil. *Appl. Environ. Microbiol.* **55**:1895–1900.
 159. Long, S. R. 1989. *Rhizobium*-legume nodulation: life together in the underground. *Cell* **56**:203–214.
 160. Long, S. R., R. S. Fisher, J. Ogawa, J. Swanson, D. W. Ehrhardt, E. M. Atkinson, and J. S. Schwedock. 1991. *Rhizobium meliloti* nodulation gene regulation and molecular signals, p. 127–133. In H. Hennecke and D. P. S. Verma (ed.), *Advances in molecular genetics of plant-microbe interactions*, vol. 1. Kluwer Academic Publishers, Dordrecht, The Netherlands.
 161. Luka, S., J. Sanjuan, R. W. Carlson, and G. Stacey. 1993. *nolMNO* genes of *Bradyrhizobium japonicum* are co-transcribed with *nodYABCSUU*, and *nolO* is involved in the synthesis of the lipo-oligosaccharide nodulation signals. *J. Biol. Chem.* **268**:27053–27059.
 162. Maillet, F., F. Debellé, and J. Dénarié. 1990. Role of the *nodD* and *sydM* genes in the activation of the regulatory gene *nodD3*, and the common and host-specific *nod* genes of *Rhizobium meliloti*. *Mol. Microbiol.* **4**:1975–1984.
 163. Malek, W. 1992. The role of motility in the efficiency of nodulation by *Rhizobium meliloti*. *Arch. Microbiol.* **158**:26–28.
 164. Marie, C., M.-A. Barny, and J. A. Downie. 1992. *Rhizobium leguminosarum* has two glucosamine synthases, *GlmS* and *NodM*, required for nodulation and development of nitrogen fixing nodules. *Mol. Microbiol.* **6**:843–851.
 165. Martinez, E., R. Puopot, J. C. Promé, M. A. Pardo, L. Segovia, G. Truchet, and J. Dénarié. 1993. Chemical signaling of *Rhizobium* nodulating bean, p. 171–176. In R. Palacios, J. Mora, and W. E. Newton (ed.), *New horizons in nitrogen fixation*. Kluwer Academic Publishers, Dordrecht, The Netherlands.
 166. Martinez, E., D. Romero, and R. Palacios. 1990. The *Rhizobium* genome. *Crit. Rev. Plant Sci.* **9**:59–93.
 167. Martinez-Romero, E. 1994. Recent development in *Rhizobium* taxonomy. *Plant Soil* **161**:11–20.
 168. Maxwell, C. A., and D. A. Phillips. 1990. Concurrent synthesis and release of *nod*-gene-inducing flavonoids from alfalfa roots. *Plant Physiol.* **93**:1552–1558.
 169. McIver, J., M. A. Djordjevic, J. J. Weinman, G. L. Bender, and B. G. Rolfe. 1989. Extension of host range of *Rhizobium leguminosarum* bv. *trifolii* caused by point mutations in *nodD* that result in alterations in regulatory function and recognition of inducer molecules. *Mol. Plant-Microbe Interact.* **2**:97–106.
 170. Meinhardt, L. W., H. B. Krishnan, P. A. Balatti, and S. G. Pueppke. 1993. Molecular cloning and characterization of a sym plasmid locus that regulates cultivar-specific nodulation of soybean by *Rhizobium fredii* USDA257. *Mol. Microbiol.* **9**:17–29.
 171. Mellor, H. Y., A. R. Glenn, R. Arwas, and M. J. Dilworth. 1987. Symbiotic and competitive properties of motility mutants of *Rhizobium trifolii* TA1. *Arch. Microbiol.* **148**:34–39.
 172. Mergaert, P., M. Van Montagu, J.-C. Promé, and M. Holsters. 1993. Three unusual modifications, a D-arabinosyl, a N-methyl, and a carbamoyl group, are present on the Nod factors of *Azorhizobium caulinodans* strain ORS571. *Proc. Natl. Acad. Sci. USA* **90**:1551–1555.
 173. Michiels, J., P. De Wilde, and J. Vanderleyden. 1993. Cloning and sequence of the *Rhizobium leguminosarum* biovar *phaseoli* *sydM* gene. *Nucleic Acids Res.* **21**:3893.
 174. Mills, K. M., and W. D. Bauer. 1985. *Rhizobium* attachment to clover roots. *J. Cell Sci. Suppl.* **2**:333–345.
 175. Mulligan, J. T., and S. R. Long. 1985. Induction of *Rhizobium meliloti* *nodC* expression by plant exudate requires *nodD*. *Proc. Natl. Acad. Sci. USA* **82**:6609–6613.
 176. Mulligan, J. T., and S. R. Long. 1989. A family of activator genes regulates expression of *Rhizobium meliloti* nodulation genes. *Genetics* **122**:7–18.
 177. Nap, J. P., and T. Bisseling. 1990. Developmental biology of a plant-prokaryote symbiosis: the legume root nodule. *Science* **250**:948–954.
 178. Ndoye, I., F. de Billy, J. Vasse, B. Dreyfus, and G. Truchet. 1994. Root nodulation of *Sesbania rostrata*. *J. Bacteriol.* **176**:1060–1068.
 179. Newcomb, W., D. Spippell, and R. L. Peterson. 1979. The early morphogenesis of *Glycine max* and *Pisum sativum* root nodules. *Can. J. Bot.* **57**:2603–2616.
 180. Newcomb, W. 1976. A correlated light electron microscopic study of symbiotic growth and differentiation in *Pisum sativum* root hair nodules. *Can. J. Bot.* **54**:2163–2186.
 181. Newcomb, W. 1981. Nodule morphogenesis and differentiation. *Int. Rev. Cytol. Suppl.* **13**:246–298.
 182. Nieuwkoop, A. J., Z. Banfalvi, N. Deshmene, D. Gerhold, M. G. Schell, K. M. Sirotkin, and G. Stacey. 1987. A locus encoding host range is linked to the common nodulation genes of *Bradyrhizobium japonicum*. *J. Bacteriol.* **169**:2631–2638.

183. Noel, K. D. 1992. Rhizobial polysaccharides required in symbiosis with legumes, p. 341–358. In D. P. S. Verma (ed.), *Molecular signals in plant-microbe communication*. CRC Press, Inc., Boca Raton, Fla.
184. Pado, C. O., and R. T. Sauer. 1984. Protein-DNA recognition. *Annu. Rev. Biochem.* **53**:293–321.
185. Peters, N. K., J. W. Frost, and S. R. Long. 1986. A plant flavone, luteolin, induces expression of *Rhizobium meliloti* nodulation genes. *Science* **233**: 977–980.
186. Peters, N. K., and S. R. Long. 1988. Alfalfa root exudates and compounds which promote or inhibit induction of *Rhizobium meliloti* nodulation genes. *Plant Physiol.* **88**:396–400.
187. Peters, N. K., and D. P. S. Verma. 1990. Phenolic compounds as regulators of gene expression in plant-microbe interactions. *Mol. Plant-Microbe Interact.* **5**:33–37.
188. Phillips, D. A. 1992. Flavonoids: plant signals to soil microbes. *Recent Adv. Phytochem.* **26**:201–231.
189. Phillips, D. A., C. M. Joseph, and C. A. Maxwell. 1992. Trigonelline and stachydrine from alfalfa seeds activate NodD2 protein in *Rhizobium meliloti*. *Plant Physiol.* **99**:1526–1531.
190. Polhill, R. M., and P. H. Raven. 1981. *Advances in legume systematics*. Royal Botanic Gardens, Kew, England.
191. Poupot, R., E. Martinez-Romero, and J. C. Promé. 1993. Nodulation factors from *Rhizobium tropici* are sulfated or nonsulfated chitopentasaccharides containing an N-methyl-N-acylglucosaminyl terminus. *Biochemistry* **32**:10430–10435.
192. Price, N. P., B. Relic, F. Talmont, A. Lewin, D. Promé, S. G. Pueppke, F. Maillet, J. Dénarié, J.-C. Promé, and W. J. Broughton. 1992. Broad-host-range *Rhizobium* species strain NGR234 secretes a family of carbamoylated, and fucosylated, nodulation signals that are O-acetylated or sulphated. *Mol. Microbiol.* **6**:3575–3584.
193. Pueppke, S. G. 1984. Absorption of slow- and fast-growing rhizobia to soybean and cowpea roots. *Plant Physiol.* **75**:924–928.
194. Rao, R. V., and D. L. Keister. 1978. Infection threads in root-hairs of soybean (*Glycine max*) plants inoculated with *Rhizobium japonicum*. *Protoplasma* **97**:311–316.
195. Recourt, K., J. Schripsema, J. W. Kijne, A. A. N. van Brussel, and B. J. J. Lugtenberg. 1991. Inoculation of *Vicia sativa* subsp. *nigra* roots with *Rhizobium leguminosarum* biovar *viciae* results in release of nod gene activating flavonones and chalcones. *Plant Mol. Biol.* **16**:841–852.
196. Recourt, K., A. A. N. van Brussel, A. J. M. Driessen, and B. J. J. Lugtenberg. 1989. Accumulation of a nod gene inducer, the flavonoid naringenin, in the cytoplasmic membrane of *Rhizobium leguminosarum* biovar *viciae* is caused by the pH-dependent hydrophobicity of naringenin. *J. Bacteriol.* **171**:4370–4377.
197. Redmond, J. W., M. Batley, M. A. Djordjevic, R. W. Innes, P. L. Kuempel, and B. G. Rolfe. 1986. Flavones induce expression of nodulation genes in *Rhizobium*. *Nature (London)* **323**:632–635.
198. Relic, B., R. Fellay, A. Lewin, X. Perret, N. P. J. Price, P. Rochepeau, and W. J. Broughton. 1993. nod genes and Nod factors of *Rhizobium* species NGR234, p. 183–190. In R. Palacios, J. Mora, and W. E. Newton (ed.), *New horizons in nitrogen fixation*. Kluwer Academic Publishers, Dordrecht, The Netherlands.
199. Relic, B., X. Perret, M. T. Estrada-Garcia, J. Kopcinska, W. Golinowski, H. B. Krishnan, S. G. Pueppke, and W. J. Broughton. 1994. Nod factors of *Rhizobium* are a key to the legume door. *Mol. Microbiol.* **13**:171–178.
200. Relic, B., F. Talmont, J. Kopcinska, W. Golinowski, J. C. Promé, and W. J. Broughton. 1993. Biological activity of *Rhizobium* sp. NGR234 Nod-factors on *Macroptilium atropurpureum*. *Mol. Plant-Microbe Interact.* **6**:764–774.
201. Reuber, T. L., S. Long, and G. C. Walker. 1991. Regulation of *Rhizobium meliloti* exo genes in free-living cells and in planta examined by using Tnpho fusions. *J. Bacteriol.* **173**:426–434.
202. Reuber, T. L., J. W. Reed, J. Glazebrook, A. Urzainqui, and G. C. Walker. 1991. Analysis of the roles of *R. meliloti* exopolysaccharides in nodulation, p. 182–188. In H. Hennecke and D. P. S. Verma (ed.), *Advances in molecular genetics of plant-microbe interactions*. Kluwer Academic Publishers, Dordrecht, The Netherlands.
203. Reuhs, B. L., R. W. Carlson, and J. S. Kim. 1993. *Rhizobium fredii* and *Rhizobium meliloti* produce 3-deoxy-D-mannose-2-oxulosonic acid containing polysaccharides that are structurally analogous to group-II-K-antigens (capsular polysaccharides) found in *Escherichia coli*. *J. Bacteriol.* **175**:3570–3580.
204. Roche, P., F. Debelle, F. Maillet, P. Lerouge, C. Faucher, G. Truchet, J. Dénarié, and J.-C. Promé. 1991. Molecular basis of symbiotic host specificity in *Rhizobium meliloti*: nodH and nodPQ genes encode the sulfation of lipo-oligosaccharide signals. *Cell* **67**:1131–1143.
205. Röhrig, H., J. Schmidt, U. Wieneke, E. Kondorosi, I. Barlier, J. Schell, and M. John. 1994. Biosynthesis of lipooligosaccharide nodulation factors—*Rhizobium* NodA protein is involved in N-acylation of the chito-oligosaccharide backbone. *Proc. Natl. Acad. Sci. USA* **91**:3122–3126.
206. Rolfe, B., and P. M. Gresshoff. 1988. Genetic analysis of legume nodule initiation. *Annu. Rev. Plant Physiol.* **39**:297–319.
207. Rolfe, B. G., M. Batley, J. W. Redmond, A. E. Richardson, R. J. Simpson, B. J. Bassam, C. L. Sargent, J. J. Weinman, M. A. Djordjevic, and F. B. Dazzo. 1988. Phenolic compounds secreted by legumes, p. 405–409. In H. Bothe, F. J. de Bruijn, and W. E. Newton (ed.), *Nitrogen fixation: hundred years after*. Gustav Fisher Verlag, Stuttgart, Germany.
208. Rossen, L., C. A. Shearman, A. W. B. Johnston, and J. A. Downie. 1985. The nodD gene of *Rhizobium leguminosarum* is autoregulatory and in the presence of plant exudate induces the nodA, B, C genes. *EMBO J.* **4**:3369–3373.
209. Rostas, K., E. Kondorosi, B. Horvath, A. Simoncsits, and A. Kondorosi. 1986. Conservation of extended promoter regions of nodulation genes in *Rhizobium*. *Proc. Natl. Acad. Sci. USA* **83**:1757–1761.
210. Rushing, B. G., M. M. Yelton, and S. R. Long. 1991. Genetic and physical analysis of the nodD3 region of *Rhizobium meliloti*. *Nucleic Acids Res.* **19**:921–927.
211. Sadowsky, M. J., P. B. Cregan, M. Gottfert, A. Sharma, D. Gerhold, F. Rodriguez-Quinones, H. H. Keyser, H. Hennecke, and G. Stacey. 1991. The *Bradyrhizobium japonicum* nola gene and its involvement in the genotype-specific nodulation of soybeans. *Proc. Natl. Acad. Sci. USA* **88**:637–641.
212. Sanjuan, J., R. W. Carlson, H. P. Spaink, U. R. Bhat, W. M. Barbour, J. Glushka, and G. Stacey. 1992. A 2-O-methylfucose moiety is present in the lipo-oligosaccharide nodulation signal of *Bradyrhizobium japonicum*. *Proc. Natl. Acad. Sci. USA* **89**:8789–8793.
213. Sanjuan, J., T. C. Dockendorff, S. Luka, and G. Stacey. 1992. Transcriptional regulation of *Bradyrhizobium japonicum* nodulation genes, p. 125. In *Abstracts of the 6th International Symposium on Molecular Plant-Microbe Interactions*.
214. Sanjuan, J., P. Grob, M. Gottfert, H. Hennecke, and G. Stacey. 1994. nodW is essential for full expression of the common nodulation genes in *Bradyrhizobium japonicum*. *Mol. Plant-Microbe Interact.* **7**:364–369.
215. Scheres, B., C. van de Wiel, A. Zalsensky, B. Horvath, H. P. Spaink, H. van Eck, F. Zwartkruis, A.-M. Wolters, T. Gloudemans, A. van Kammen, and T. Bisseling. 1990. The ENOD12 gene product is involved in the infection process during the pea-*Rhizobium* interaction. *Cell* **60**:281–294.
216. Scheu, A. K., A. Economou, G. F. Hong, S. Ghelani, A. W. B. Johnston, and J. A. Downie. 1992. Secretion of the *Rhizobium leguminosarum* nodulation protein NodO by haemolysin-type systems. *Mol. Microbiol.* **6**:231–238.
217. Schlaman, H. R. M. 1992. Regulation of nodulation gene expression in *Rhizobium leguminosarum* biovar *viciae*. Ph.D. thesis, University of Leiden, Leiden, The Netherlands.
218. Schlaman, H. R. M., B. Horvath, E. Vijgenboom, R. J. H. Okker, and B. J. J. Lugtenberg. 1991. Suppression of nodulation gene expression in bacteroids of *Rhizobium leguminosarum* bv. *viciae*. *J. Bacteriol.* **173**:4277–4287.
219. Schlaman, H. R. M., B. J. J. Lugtenberg, and R. J. H. Okker. 1992. The NodD protein does not bind to the promoter of inducible nodulation genes in bacteroids of *Rhizobium leguminosarum* bv. *viciae*. *J. Bacteriol.* **174**:6109–6116.
220. Schlaman, H. R. M., R. J. H. Okker, and B. J. J. Lugtenberg. 1990. Subcellular localization of the *Rhizobium leguminosarum* nodI gene product. *J. Bacteriol.* **172**:5486–5498.
221. Schlaman, H. R. M., R. J. H. Okker, and B. J. J. Lugtenberg. 1992. Regulation of nodulation gene expression by NodD in rhizobia. *J. Bacteriol.* **174**:5177–5182.
222. Schlaman, H. R. M., H. P. Spaink, R. J. H. Okker, and B. J. J. Lugtenberg. 1989. Subcellular localization of the nodD gene product in *Rhizobium leguminosarum*. *J. Bacteriol.* **171**:4648–4693.
223. Schmidt, P. E., W. J. Broughton, and D. Werner. 1994. Nod factor of *Bradyrhizobium japonicum* and *Rhizobium* sp. NGR234 induce flavonoid accumulation in soybean root exudate. *Mol. Plant-Microbe Interact.* **7**:384–390.
224. Schofield, P. R., and J. M. Watson. 1986. DNA sequence of *Rhizobium trifolii* nodulation genes reveals a reiterated and potentially regulatory sequence preceding nodABC and nodEF. *Nucleic Acids Res.* **14**:2891–2903.
225. Schultze, M., E. Kondorosi, A. Kondorosi, C. Staehelin, R. Mellor, and T. Boller. 1993. The sulfate group on the reducing end protects Nod signals of *Rhizobium meliloti* against hydrolysis by *Medicago* chitinases, p. 159–164. In R. Palacios, J. Mora, and W. E. Newton (ed.), *New horizons in nitrogen fixation*. Kluwer Academic Publishers, Dordrecht, The Netherlands.
226. Schultze, M., B. Quiclet-Sire, E. Kondorosi, H. Virelizier, J. N. Glushka, G. Endre, S. D. Géro, and A. Kondorosi. 1992. *Rhizobium meliloti* produces a family of sulfated lipo-oligosaccharides exhibiting different degrees of plant host specificity. *Proc. Natl. Acad. Sci. USA* **89**:192–196.
227. Schwedock, J., and S. R. Long. 1990. ATP sulphurylase activity of the NodP and NodQ gene products of *Rhizobium meliloti*. *Nature (London)* **348**:644–647.
228. Schwedock, J., and S. R. Long. 1989. Nucleotide sequence and protein products of two new nodulation genes of *Rhizobium meliloti*, nodP and nodQ. *Mol. Plant-Microbe Interact.* **2**:181–194.
229. Scott, R. F. 1986. Conserved nodulation genes from the non-legume *Bradyrhizobium* sp. (Parasponia). *Nucleic Acids Res.* **14**:2905–2910.
230. Sharma, S. B., and E. R. Signer. 1990. Temporal and spatial regulation of the symbiotic genes of *Rhizobium meliloti* in planta revealed by transposon Tn5-gusA. *Genes Dev.* **4**:344–356.
231. Shearman, C. A., L. Rossen, A. W. B. Johnston, and J. A. Downie. 1986. The

- Rhizobium leguminosarum* nodulation gene *nodF* encodes a polypeptide similar to acyl-carrier protein and is regulated by *nodD* plus a factor in pea root exudate. EMBO J. 5:647-652.
232. Sheldon, P. S., R. G. O. Kekwick, C. Sidebottom, C. G. Smith, and A. R. Slabas. 1990. 3-Oxoacyl-(acyl-carrier-protein) reductase from avocado (*Persea americana*) fruit mesocarp. Biochem. J. 271:713-720.
 233. Smit, G., J. W. Kijne, and B. J. J. Lugtenberg. 1986. Correlation between extracellular fibrils and attachment of *Rhizobium leguminosarum* to pea root hair tips. J. Bacteriol. 168:821-827.
 234. Smit, G., J. W. Kijne, and B. J. J. Lugtenberg. 1987. Involvement of both cellulose fibrils and a Ca^{2+} -dependent adhesin in the attachment of *Rhizobium leguminosarum* to pea root hair tips. J. Bacteriol. 169:4294-4301.
 235. Smit, G., J. W. Kijne, and B. J. J. Lugtenberg. 1989. Roles of flagella, lipopolysaccharide, and a Ca^{2+} -dependent cell surface protein in attachment of *Rhizobium leguminosarum* biovar *viciae* to pea root hair tips. J. Bacteriol. 171:569-572.
 236. Smit, G., T. J. J. Logman, M. E. T. I. Boerrigter, J. W. Kijne, and B. J. J. Lugtenberg. 1989. Purification and partial characterization of the Ca^{2+} -dependent adhesin from *Rhizobium leguminosarum* biovar *viciae*, which mediates the first step in attachment of *Rhizobiaceae* cells to plant root hair tips. J. Bacteriol. 171:4054-4062.
 237. Smit, G., T. J. J. Logman, M. E. T. I. Boerrigter, J. W. Kijne, and B. J. J. Lugtenberg. 1989. Purification and partial characterization of the *Rhizobium leguminosarum* bv. *viciae* Ca^{2+} dependent adhesin with mediates the first step in attachment of all of the family *Rhizobiaceae* to plant root hair tips. J. Bacteriol. 171:4054-4062.
 238. Smit, G., S. Swart, B. J. J. Lugtenberg, and J. Kijne. 1992. Molecular mechanisms of attachment of *Rhizobium* bacteria to plant roots. Mol. Microbiol. 6:2897-2903.
 239. Smit, G., D. M. J. Tubbing, J. W. Kijne, and B. J. J. Lugtenberg. 1991. Role of Ca^{2+} in the activity of rhicadhesin from *Rhizobium leguminosarum* biovar *viciae*, which mediates the first step in attachment of *Rhizobiaceae* cells to plant root hair tips. Arch. Microbiol. 155:278-283.
 240. Spaink, H. P. 1992. Rhizobial lipo-oligosaccharides: answers and questions. Plant Mol. Biol. 20:977-987.
 241. Spaink, H. P., A. Aarts, G. V. Bloemberg, J. Folch, O. Geiger, H. R. M. Schlaman, J. E. Thomas-Oates, A. A. N. van Brussel, K. van de Sande, P. van Spronsen, A. H. M. Wijffes, and B. J. J. Lugtenberg. 1992. Rhizobial lipo-oligosaccharide signals: their biosynthesis and their role in the plant, p. 151-162. In E. W. Nester and D. P. S. Verma (ed.), Advances in molecular genetics of plant-microbe interactions, vol. 2. Kluwer Academic Publishers, Dordrecht, The Netherlands.
 242. Spaink, H. P., A. Aarts, G. Stacey, G. V. Bloemberg, B. J. J. Lugtenberg, and E. P. Kennedy. 1992. Detection and separation of *Rhizobium* and *Bradyrhizobium* Nod metabolites using thin layer chromatography. Mol. Plant-Microbe Interact. 5:72-80.
 243. Spaink, H. P., R. J. H. Okker, C. A. Wijffelman, E. Pees, and B. J. J. Lugtenberg. 1987. Promoters in the nodulation region of the *Rhizobium leguminosarum* Sym plasmid pRL1J1. Plant Mol. Biol. 9:27-39.
 244. Spaink, H. P., R. J. H. Okker, C. A. Wijffelman, T. Tak, L. Goossen-de Roo, E. Pees, A. A. N. van Brussel, and B. J. J. Lugtenberg. 1989. Symbiotic properties of rhizobia containing a flavonoid-independent hybrid *nodD* product. J. Bacteriol. 171:4045-4053.
 245. Spaink, H. P., D. M. Sheeley, A. A. N. van Brussel, J. Glushka, W. S. York, T. Tak, O. Geiger, E. P. Kennedy, V. N. Reinhold, and B. J. J. Lugtenberg. 1991. A novel highly unsaturated fatty acid moiety of lipo-oligosaccharide signals determines host specificity of *Rhizobium*. Nature (London) 354:125-130.
 246. Spaink, H. P., J. Weinman, M. A. Djordjevic, C. A. Wijffelman, J. H. Okker, and B. J. J. Lugtenberg. 1989. Genetic analysis and cellular localization of the *Rhizobium* host specificity-determining NodE protein. EMBO J. 8:2811-2818.
 247. Spaink, H. P., C. A. Wijffelman, E. Pees, R. J. H. Okker, and B. J. J. Lugtenberg. 1987. *Rhizobium* nodulation gene *nodD* as a determinant of host specificity. Nature (London) 328:337-340.
 248. Spaink, H. P., C. A. Wijffelman, E. Pees, R. J. H. Okker, and B. J. J. Lugtenberg. 1989. Localization and functional regions of the *Rhizobium nodD* product using hybrid *nodD* genes. Plant Mol. Biol. 12:59-73.
 249. Spaink, H. P., A. H. M. Wijffes, O. Geiger, G. V. Bloemberg, T. Ritsema, and B. J. J. Lugtenberg. 1993. The function of the rhizobial *nodABC* and *nodFEL* operons in the biosynthesis of lipo-oligosaccharides, p. 165-170. In R. Palacios, J. Mora, and W. E. Newton (ed.), New horizons in nitrogen fixation. Kluwer Academic Publishers, Dordrecht, The Netherlands.
 250. Sprent, J. L., and S. M. de Faria. 1988. Mechanisms of infection of plants by nitrogen fixing organisms. Plant Soil 110:157-165.
 251. Stacey, G. 1990. Workshop summary: compilation of the *nod*, *fix* and *nif* genes of rhizobia and information concerning their function, p. 239-244. In P. M. Gresshoff, L. E. Roth, G. Stacey, and W. E. Newton (ed.), Nitrogen fixation: achievements and objectives. Chapman & Hall, New York.
 252. Stacey, G., S. Luka, J. Sanjuan, Z. Banfalvi, A. J. Nieuwkoop, J. Y. Chun, L. S. Forsberg, and R. Carlson. 1994. *nodZ*, a unique host-specific nodulation gene, is involved in the fucosylation of the lipooligosaccharide signal of *Bradyrhizobium japonicum*. J. Bacteriol. 176:620-633.
 253. Staehelin, C., M. Schultze, E. Kondorosi, R. B. Mellor, T. Boller, and A. Kondorosi. 1994. Structural modifications in *Rhizobium meliloti* Nod factors influence their stability against hydrolysis by root chitinases. Plant J. 5:319-330.
 254. Stokkermans, T. J. W., and N. K. Peters. 1994. *Bradyrhizobium elkanii* lipo-oligosaccharide signals induce complete nodule structures on *Glycine soja* Siebold et Zucc. Planta 193:413-420.
 255. Surin, B. P., and J. A. Downie. 1988. Characterization of the *Rhizobium leguminosarum* genes *nodLMN* involved in efficient host specific nodulation. Mol. Microbiol. 2:173-183.
 256. Surin, B. P., and J. A. Downie. 1989. *Rhizobium leguminosarum* genes required for expression and transfer of host specific nodulation. Plant Mol. Biol. 12:19-29.
 257. Surin, B. P., J. M. Watson, W. D. O. Hamilton, A. Economou, and J. A. Downie. 1990. Molecular characterization of the nodulation gene, *nodT*, from two biovars of *Rhizobium leguminosarum*. Mol. Microbiol. 4:245-252.
 258. Sutton, M. J., E. J. A. Lea, and J. A. Downie. 1994. The nodulation-signaling protein NodO from *Rhizobium leguminosarum* biovar *viciae* forms ion channels in membranes. Proc. Natl. Acad. Sci. USA 91:9990-9994.
 259. Swanson, J. A., J. T. Mulligan, and S. R. Long. 1993. Regulation of *syrM* and *nodD3* in *Rhizobium meliloti*. Genetics 134:435-444.
 260. Trinick, M. J. 1979. Structure of nitrogen fixing nodules formed by *Rhizobium* on roots of *Parasponia andersonii* Planck. Can. J. Microbiol. 25:565-578.
 261. Trinick, M. J. 1988. Biology of the *Parasponia-Bradyrhizobium* symbiosis. Plant Soil 110:177-185.
 262. Trinick, M. J., and J. Galbraith. 1976. Structure of root nodules formed by *Rhizobium* on the non-legume *Trema cannabina* var *scabra*. Arch. Microbiol. 108:159-166.
 263. Truchet, G., P. Roche, P. Lerouge, J. Vasse, S. Camut, F. de Billy, J.-C. Promé, and J. Dénarié. 1991. Sulphated lipo-oligosaccharide signals of *R. meliloti* elicit root nodule organogenesis in alfalfa. Nature (London) 351:670-673.
 264. Tsien, H. C., B. L. Dreyfus, and E. L. Schmidt. 1983. Initial stages in the morphogenesis of nitrogen-fixing stem nodules of *Sesbania rostrata*. J. Bacteriol. 156:888-897.
 265. Turgeon, B. G., and W. D. Bauer. 1982. Early events in the infection of the soybean by *Rhizobium japonicum*. Time course cytology of the initial infection process. Can. J. Bot. 60:152-161.
 266. Turgeon, B. G., and W. D. Bauer. 1985. Ultrastructure of infection thread development during the infection of soybean by *Rhizobium japonicum*. Planta 163:328-349.
 267. van Brussel, A. A. N., R. Bakhuizen, P. van Spronsen, H. P. Spaink, T. Tak, B. J. J. Lugtenberg, and J. Kijne. 1992. Induction of pre-infection thread structures in the host plant by lipo-oligosaccharides of *Rhizobium*. Science 257:70-72.
 268. van Brussel, A. A. N., K. Recourt, E. Pees, H. P. Spaink, T. Tak, C. A. Wijffelman, J. W. Kijne, and B. J. J. Lugtenberg. 1990. A biovar-specific signal of *Rhizobium leguminosarum* bv. *viciae* induces increased nodulation gene-inducing activity in root exudate of *Vicia sativa* subsp. *nigra*. J. Bacteriol. 172:5394-5401.
 269. van Brussel, A. A. N., S. A. J. Zaat, H. C. J. Canter-Cremers, C. A. Wijffelman, E. Pees, and B. J. J. Lugtenberg. 1986. Role of plant root exudate and Sym plasmid-localized nodulation genes in the synthesis by *Rhizobium leguminosarum* of Trs factor, which causes thick and short roots on common vetch. J. Bacteriol. 165:517-522.
 270. van Rhijn, P. J. S., J. Desair, K. Vlassak, and J. Vanderleyden. 1994. Functional analysis of *nodD* genes of *Rhizobium tropici* CIAT899. Mol. Plant-Microbe Interact. 7:666-676.
 271. van Rhijn, P. J. S., J. Desair, K. Vlassak, and J. Vanderleyden. Unpublished data.
 - 271a. van Rhijn, P. J. S., B. Feys, C. Verreth, and J. Vanderleyden. 1993. Multiple copies of *nodD* in *Rhizobium tropici* CIAT899 and BR816. J. Bacteriol. 175:438-447.
 272. Vasse, J., F. de Billy, S. Camut, and G. Truchet. 1990. Correlation between ultrastructural differentiation of bacteroids and nitrogen fixation in alfalfa nodules. J. Bacteriol. 172:4296-4306.
 273. Vasse, J., and G. Truchet. 1984. The *Rhizobium*-legume symbiosis: observation of root infection by bright-field microscopy after staining with methylene blue. Planta 161:487-489.
 274. Vázquez, M., A. Davalos, A. de las Penas, F. Sanchez, and C. Quinto. 1991. Novel organization of the common nodulation genes in *Rhizobium leguminosarum* bv. *phaseoli* strains. J. Bacteriol. 173:1250-1258.
 275. Vázquez, M., O. Santana, and C. Quinto. 1993. The NodI and NodJ proteins from *Rhizobium* and *Bradyrhizobium* strains are similar to capsular polysaccharide secretion proteins from Gram-negative bacteria. Mol. Microbiol. 8:369-377.
 276. Verma, D. P. S., C. A. Hu, and M. Zhang. 1992. Root nodule development: origin and regulation of nodulin genes. Physiol. Plant. 85:253-265.
 277. Vesper, S. J., and W. D. Bauer. 1986. Role of pili (fimbriae) in attachment of *Bradyrhizobium japonicum* to soybean roots. Appl. Environ. Microbiol. 52:134-141.

278. **Vesper, S. J., and T. V. Bhuvaneswari.** 1988. Nodulation of soybean roots by an isolate of *Bradyrhizobium japonicum* with reduced firm attachment capability. *Arch. Microbiol.* **150**:15–19.
279. **Vesper, S. J., S. A. Malik, and W. D. Bauer.** 1987. Transposon mutants of *Bradyrhizobium japonicum* altered in attachment to host roots. *Appl. Environ. Microbiol.* **53**:1959–1961.
280. **Waelkens, F., T. Voets, K. Vlassak, J. Vanderleyden, and P. van Rhijn.** The *nodS* gene of *Rhizobium tropici* strain CIAT899 is necessary for nodulation on *Phaseolus vulgaris* and on *Leucaena leucocephala*. *Mol. Plant-Microbe Interact.*, in press.
281. **Wang, S.-P., and G. Stacey.** 1990. Ammonia regulation of *nod* genes in *Bradyrhizobium japonicum*. *Mol. Gen. Genet.* **223**:329–331.
282. **Wang, S.-P., and G. Stacey.** 1991. Studies of the *Bradyrhizobium japonicum nodD1* promoter: a repeated structure for the *nod* box. *J. Bacteriol.* **173**:3356–3365.
283. **Willems, A., and M. D. Collins.** 1993. Phylogenetic analysis of rhizobia and agrobacteria based on 16S rRNA gene sequences. *Int. J. Syst. Bacteriol.* **43**:305–313.
284. **Wood, S. M., and W. Newcomb.** 1989. Nodule morphogenesis: the early infection of alfalfa (*Medicago sativa*) root hairs by *Rhizobium meliloti*. *Can. J. Bot.* **67**:3108–3122.
285. **Yanagi, M., and K. Yamasato.** 1993. Phylogenetic analysis of the family Rhizobiaceae and related bacteria by sequencing of 16S rRNA gene using PCR and DNA sequencer. *FEMS Microbiol. Lett.* **107**:115–120.
286. **Yao, P. Y., and J. M. Vincent.** 1969. Host specificity in the root hair “curling factor” of *Rhizobium sp.* *Aust. J. Biol. Sci.* **22**:413–423.
287. **Young, J. P. W., H. L. Downer, and B. D. Eardly.** 1991. Phylogeny of the phototrophic *Rhizobium* strain Btail by polymerase chain reaction-based sequencing of the 16S rRNA gene segment. *J. Bacteriol.* **173**:2271–2277.
288. **Young, J. P. W., and A. W. B. Johnston.** 1989. The evolution of specificity in the legume-*Rhizobium* symbiosis. *Trends Ecol. Evol.* **4**:331–349.